

Sex-dependent toxoplasmosis-associated differences in testosterone concentration in humans

J. FLEGR^{1*}, J. LINDOVÁ^{1,2} and P. KODYM³

¹ Department of Parasitology, Faculty of Sciences, Charles University, Viničná 7, 128 44 Prague 2, Czech Republic

² Department of Anthropology, Faculty of Humanities, Charles University, Prague, Husníková 2075, 155 00, Prague 13, Czech Republic

³ National Reference Laboratory for Toxoplasmosis, National Institute of Public Health, Šrobárova 48, 100 42, Prague 10, Czech Republic

(Received 6 July 2007; revised 7 September 2007; accepted 27 November 2007; first published online 21 January 2008)

SUMMARY

Several lines of indirect evidence suggest that subjects with latent infection of the coccidian parasite *Toxoplasma gondii* have a higher concentration of testosterone than uninfected controls. Here, we searched for direct evidence of latent toxoplasmosis-associated differences in testosterone concentration among a population of 174 female and 91 male students screened for *Toxoplasma* infection. We have found *Toxoplasma*-infected men to have a higher concentration of testosterone and *Toxoplasma*-infected women to have a lower concentration of testosterone than *Toxoplasma*-free controls. The opposite direction of the testosterone shift in men compared to women can explain the observed gender specificity of behavioural shifts in *Toxoplasma*-infected subjects.

Key words: infection, behaviour, *Toxoplasma*, hormones, androgens, manipulation.

INTRODUCTION

A common protozoan parasite *Toxoplasma gondii* (worldwide prevalence varies between 20 and 70% in different countries, depending on climate, hygiene standards and eating habits) (Tenter *et al.* 2000) is known to induce behavioural and neurophysiological changes in infected human or animal hosts (Webster, 2001). Increased dopamine and testosterone levels are suspected to play an important role in the observed changes (Flegr, 2007). However, the association between toxoplasmosis and increased testosterone concentration was only postulated on the basis of indirect evidence. Infected men are taller, have a lower left hand 2D:4D ratio (Flegr *et al.* 2005) and higher perceived dominance and masculinity scores (Hodkova *et al.* 2007). Infected women have a (non-significantly) lower left hand 2D:4D ratio and are more likely to give birth to a boy than a girl (Kaňková *et al.* 2007). All these traits are known to be positively associated with testosterone concentration in humans (James, 1986, 1996; Drop *et al.* 1998; Brown *et al.* 2002; Okten *et al.* 2002; Neave *et al.* 2003; Lutchmaya *et al.* 2004). However, no studies directly comparing testosterone levels in *Toxoplasma*-infected and *Toxoplasma*-free subjects have been available to date and analogous studies in animal models are missing as well.

* Corresponding author: Department of Parasitology, Faculty of Science, Charles University, Viničná 7, CZ-128 44 Praha 2, Czech Republic. Tel: +420 221951821. Fax: +420 224919704. E-mail: flegr@cesnet.cz

Here, we searched for direct evidence of the changed testosterone levels in *Toxoplasma*-infected men and women by testing the concentration of testosterone in saliva in a population of 174 female and 91 male students screened for *Toxoplasma* infection.

MATERIALS AND METHODS

Subjects

Undergraduate biology students of the Faculty of Sciences, Charles University, Prague, were addressed during regular biology lectures and were invited to participate in the study on a voluntary basis. In total, 91 male students and 174 female students were enrolled in the study and signed an informed consent form. All participants provided 2 ml of blood for serological testing and 3 saliva samples of approximately 200 μ l (collected approximately at 09.00, 11.30 and 13.00). Meanwhile, all participants performed the same panel of psychological and ethological tests. All blood and saliva samples were stored in a freezer at -20°C until assayed. The recruitment of the study subjects and data handling practices complied with the Czech regulations currently in force.

Immunological tests for toxoplasmosis

All serological tests were carried out in the National Reference Laboratory for Toxoplasmosis, National

Institute of Public Health, Prague. Specific anti-*Toxoplasma* IgG ELISA (in-house test NRL TOXO – Pokorný *et al.* 1989); and the complement fixation test (CFT – SEVAPHARMA, Prague) which is more reliable in old *T. gondii* infections as decrease in CFT titres is more regular (Warren and Sabin, 1942; Kodým *et al.* 2007) were performed in all subjects. Anti-*Toxoplasma* IgM antibodies were determined by ELISA (TestLine, Brno) optimized for early detection of acute toxoplasmosis in subjects with CFT titres of $\geq 1:64$. *Toxoplasma* antibody titres in the sera were measured at dilutions between 1:8 and 1:1024. The subjects with positive IgG ELISA values, CFT titres higher or equal than 1:8 and testing IgM negative by ELISA (positivity index < 0.9) were considered latent toxoplasmosis positive.

Radioimmunoassay (RIA) test for testosterone

All testosterone assays were performed at the Institute of Endocrinology, Prague. Saliva samples and controls or blanks (double-distilled water), 1.0 ml each, were spiked with [^3H] testosterone (Radiochemical Centre, Amersham, UK, 1200 dpm/sample), and extracted in duplicate with diethyl ether (4 ml) in stoppered glass tubes. The aqueous phase was frozen in solid carbon dioxide, the organic phase was decanted and the ether was evaporated to dryness. The extracts were dissolved in ethanol (500 μl), and then 100 μl of the solution were removed for determination of the losses during the extraction, while the rest was evaporated again and taken for radioimmunoassay. A standard curve consisting of 0.025, 0.05, 0.1, 0.2, 0.4, 0.8, 1.6 and 3.2 nmol/l testosterone in duplicate was prepared. Antiserum (rabbit anti-testosterone-3-CMO: BSA, working dilution 1:100 000) and the tracer ([^{125}I] iodohistaminyl testosterone derivative, 15 000 cpm), 100 μl each, were added, the volume was adjusted to 300 μl with the working buffer (20 mmol sodium phosphate saline containing sodium azide and BSA, 0.1% each) and the tubes were equilibrated at room temperature for 1 h or overnight at 4 °C. After incubation, 1 ml of dextran-coated charcoal suspension (0.025 and 0.25 g/100 ml, respectively) was added to each tube to separate the free fraction and the radioactivity of ^{125}I was measured in the supernatant using a 12-channel gamma counter (Berthold, FRG). Results were calculated from the standard curve using a log-logit transformation, corrected for recovery and expressed as nmol of testosterone per litre of sample. The samples were assayed in triplicate. Because of the low volume of some saliva samples (the minimum limit for a reliable assay was 100 μl) the complete testosterone concentration data (all 3 samples) were not obtained for all subjects.

Statistics

Differences in age and testosterone concentration between men and women were measured with *t*-tests. The relation between the average testosterone concentration in all (usually 3) samples of a particular subject and independent factors sex, age and *Toxoplasma* infection was tested with General Linear Models (GLM). The difference was also analysed with repeated measures GLM, with concentration of testosterone in samples 1–3 as a dependent variable and sex, age and toxoplasmosis as independent variables, in 236 subjects with available data from all 3 saliva samples (126 *Toxoplasma*-infected and 26 *Toxoplasma*-free women, 62 *Toxoplasma*-infected and 22 *Toxoplasma*-free men). If the Mauchly's test indicated violation of the sphericity assumption, the degrees of freedom were corrected using the Greenhouse-Geisser method. The separate analyses for men and women were performed with GLM (with the covariate age) and (nonparametric) Mann-Whitney U test (without any covariate). All parametric statistical tests were performed with log-transformed testosterone concentrations; however, there was practically no qualitative difference between results of statistical tests performed either with or without log transformation of the testosterone concentration data. All analyses including the tests of analyses assumptions (Levene tests of homogeneity, univariate and multivariate Lack of fit tests, Mauchly's tests of sphericity) were performed with SPSS 12.0.

RESULTS

The average age of 174 women was 21.03 years, range 18–28, median 21, lower quartile 20, upper quartile 22, average concentration of testosterone was 0.230 ng/ml, range 0.035–0.860, median 0.201, lower quartile 0.148, upper quartile 0.275. The average age of 91 men was 20.91 years, range 19–27, median 20, lower quartile 20, upper quartile 22, average concentration of testosterone was 0.387 ng/ml, range 0.083–1.088, median 0.358, lower quartile 0.266, upper quartile 0.485. The difference in age between women and men was non-significant $t_{263} = 0.494$, $P = 0.62$. However, we found significantly higher levels of testosterone ($t_{263} = -8.133$, $P < 0.0001$) in men than in women.

Twenty-nine (16.7%) women and 23 (25.3%) men were *Toxoplasma*-infected. The difference between *Toxoplasma*-infected and *Toxoplasma*-free subjects in average testosterone concentration estimated by GLM with independent factors sex, age and toxoplasmosis was nonsignificant ($F_{1,260} = 0.906$, $P = 0.342$, $\eta^2 = 0.003$). At the same time, the effect of the toxoplasmosis-sex interaction on the testosterone concentration was highly significant ($F_{1,260} = 13.434$, $P < 0.001$, $\eta^2 = 0.049$). Infected men had higher and




Fig. 1. Average testosterone levels in saliva of *Toxoplasma*-free and *Toxoplasma*-infected men and women. Boxes represent standard errors; spreads represent standard deviations.

infected women had lower testosterone levels than *Toxoplasma*-free men and women, respectively (Fig. 1). The difference in testosterone concentration was significant for women (GLM: $F_{1,171}=12.643$, $P<0.001$, $\eta^2=0.069$, M-W: $U=1298$, $P=0.001$) and non-significant for men ($F_{1,88}=2.678$, $P=0.105$, $\eta^2=0.030$, M-W: $U=645$, $P=0.211$).

The results of repeated measures GLM, with concentration of testosterone in samples 1–3 as a dependent variable and sex, age and toxoplasmosis as independent variables, were similar to those of GLM analyses with the average testosterone concentration for samples 1–3 as a dependent variable. The effect of toxoplasmosis was non-significant ($F_{1,231}=1.155$, $P=0.284$, $\eta^2=0.005$) while the effects of toxoplasmosis-sex interaction ($F_{1,231}=16.347$, $P<0.001$, $\eta^2=0.066$) was significant. The effect of time-toxoplasmosis-sex interaction was non-significant ($F_{1,86,429-55}=2.464$, $P=0.090$, $\eta^2=0.011$, linear contrast: $F_{1,231}=0.207$, $P=0.649$, $\eta^2<0.001$, quadratic contrast: $F_{1,231}=6.047$, $P=0.015$, $\eta^2=0.026$). Separate repeated measures GLM analyses for women and men have shown significant effects of toxoplasmosis in women ($F_{1,149}=15.579$, $P<0.001$, $\eta^2=0.095$) and of time-toxoplasmosis interaction in men ($F_{2,162}=3.283$, $P<0.040$, $\eta^2=0.039$, linear contrast: $F_{1,81}=0.095$, $P=0.759$, $\eta^2=0.001$, quadratic contrast: $F_{1,81}=7.959$, $P=0.006$, $\eta^2=0.089$), (Fig. 2).

DISCUSSION

We have found *Toxoplasma*-infected men to have a higher concentration of testosterone and *Toxoplasma*-infected women to have a lower concentration of testosterone than *Toxoplasma*-free subjects. The difference was significant for all 3 saliva samples in women but only for saliva samples 1 (collected at 09.00) and 3 (collected at 13.00) in men.

The proximal mechanism of association between testosterone and toxoplasmosis is not clear. As




Fig. 2. Changes of testosterone level in saliva of *Toxoplasma*-free (empty circles) and *Toxoplasma*-infected (black circles) subjects between 09.00 and 13.00. Vertical bars denote 0.95 confidence intervals.

stressed in previous articles (Flegr *et al.* 2005; Hodkova *et al.* 2007), a case control study cannot decide whether *Toxoplasma* infection induces changes in testosterone concentration or whether low-testosterone and high-testosterone subjects differ in the probability of acquiring *Toxoplasma* infection or both. High concentrations of testosterone are known to have immunosuppressive effects (Roberts *et al.* 2001; Schuster and Schaub, 2001). The results of the present study, in particular the lower concentration of testosterone in *Toxoplasma* infected women, however, make the immunosuppression-based explanation of the association between *Toxoplasma* infection and testosterone concentration rather unlikely.

The increased level of testosterone in men could be explained by a positive association between testosterone and dopamine (Szczycka *et al.* 1998; Dominguez and Hull, 2005; Hull *et al.* 2004) that can increase in response to local inflammatory processes in the infected brain (Flegr, 2007). However, we have no explanation for the decreased testosterone in infected women.

It has been shown in several studies that latent toxoplasmosis has gender-different effects on the personality factors A (warmth), G (rule consciousness), L (vigilance, mistrust) and Q3 (self-control, self-image) from Cattell's 16PF Questionnaire (Flegr *et al.* 1996, 2000). Behavioural experiments also showed gender-different effects in behavioural variables Self-Control and Clothes Tidiness (analogue to the 16PF factors G – conscientiousness and Q3 – self-control), with infected men scoring significantly lower than uninfected men and infected women scoring non-significantly higher than uninfected women (Lindová *et al.* 2006).

Biological and psychological hypotheses were suggested to explain the opposite directions of personality factor scores and behavioural shifts in men

and women. According to the biological hypothesis, infected intermediate hosts are manipulated by the parasite to behave in a way which makes them easier prey for the final host (Lindová *et al.* 2006). In primitive human societies, being reserved, detached and critical of other people, being uncontrolled, lax, disregarding rules (low Self Control) and untidily dressed would isolate males from a group, making them a more probable prey. The opposite traits in infected woman could make her more attractive for potential sexual partners, and in such a way increase the rate of its transmission into the offspring.

The psychological hypothesis assumes that men and women use a different strategy to cope with non-specific stressors (including toxoplasmosis) (Lindová *et al.* 2006). There is some evidence that men are socialized to cope with stress differently from women. Men are expected to use more problem-focused forms of coping, while women are brought up to cope with an emotion-focused style (Brems and Johnson, 1989; Carver *et al.* 1989). Thus, coping with toxoplasmosis, men would be expected to withdraw from society in order to concentrate on the 'problem', whereas women would be expected to turn to society, where they can express their emotions.

Our present results suggest that the gender-different effects of latent toxoplasmosis on behavioural traits and personality factor scores can be the product of the opposite effects of toxoplasmosis on testosterone levels. The changes in testosterone concentration in men and women can be either side-effects of the *Toxoplasma* activities or a result of the specific manipulation activity of the parasite aimed to increase the probability of its transmission from intermediate to definitive host (Dawkins, 1982; Barnard and Behnke, 1990). Therefore, our testosterone hypothesis is compatible with the biological explanation mentioned above.

Although several pieces of indirect evidence for increased testosterone in *Toxoplasma*-infected men have been recently reported, we present here the first piece of direct (biochemical) evidence for the changed testosterone concentration in saliva of infected subjects. Surprisingly, our study has not only confirmed a positive association between latent toxoplasmosis and testosterone level in infected men but also has shown a negative association between latent toxoplasmosis and testosterone level in infected women. Obviously, it is not possible to perform an infection experiment in the human-*Toxoplasma* system. The relatively low incidence of toxoplasmosis would also make a longitudinal study rather inefficient. Therefore, the direction of causality between toxoplasmosis and testosterone level will probably need to be studied in experimentally infected laboratory animals.

We thank Aleš Kuběna for his help with statistical analysis. This research was supported by grant no. 406/07/0581 of

the Grant Agency of the Czech Republic, 160/2006/B-B10/PrF GAUK and grant no. 0021620828 of the Czech Ministry of Education.

REFERENCES

- Barnard, C. J. and Behnke, J. M.** (1990). *Parasitism and Host Behaviour*. 1st Edn. Taylor and Francis, New York.
- Brems, C. and Johnson, M. E.** (1989). Problem-solving appraisal and coping style – the influence of sex-role orientation and gender. *Journal of Psychology* **123**, 187–194.
- Brown, W. M., Hines, M., Fane, B. A. and Breedlove, S. M.** (2002). Masculinized finger length patterns in human males and females with congenital adrenal hyperplasia. *Hormones and Behavior* **42**, 380–386.
- Carver, C. S., Scheier, M. F. and Weintraub, J. K.** (1989). Assessing coping strategies – a theoretically based approach. *Journal of Personality and Social Psychology* **56**, 267–283.
- Dawkins, R.** (1982). *The Extended Phenotype, the Gene as the Unit of Selection*. 1st Edn. W.H. Freeman, Oxford.
- Drop, S. L. S., De Waal, W. J. and Keizer-Schrama, S. M. P. F.** (1998). Sex steroid treatment of constitutionally tall stature. *Endocrine Reviews* **19**, 540–558.
- Dominguez, J. M. and Hull, E. M.** (2005). Dopamine, the medial preoptic area, and male sexual behavior. *Physiology and Behavior* **86**, 356–368.
- Flegr, J.** (2007). Effects of *Toxoplasma* on human behavior. *Schizophrenia Bulletin* **33**, 757–760.
- Flegr, J., Hruskova, M., Hodny, Z., Novotna, M. and Hanusova, J.** (2005). Body height, body mass index, waist-hip ratio, fluctuating asymmetry and second to fourth digit ratio in subjects with latent toxoplasmosis. *Parasitology* **130**, 621–628.
- Flegr, J., Kodym, P. and Tolarova, V.** (2000). Correlation of duration of latent *Toxoplasma gondii* infection with personality changes in women. *Biological Psychology* **53**, 57–68.
- Flegr, J., Zitkova, S., Kodym, P. and Frynta, D.** (1996). Induction of changes in human behaviour by the parasitic protozoan *Toxoplasma gondii*. *Parasitology* **113**, 49–54.
- Hodkova, H., Kolbekova, P., Skallova, A., Lindová, J. and Flegr, J.** (2007). Higher perceived dominance in *Toxoplasma* infected men – the new evidence for role of increased level of testosterone in toxoplasmosis-associated changes in human behavior. *Neuroendocrinology Letters* **28**, 101–105.
- Hull, E. M., Muschamp, J. W. and Sato, S.** (2004). Dopamine and serotonin: influences on male sexual behavior. *Physiology and Behavior* **83**, 291–307.
- James, W. H.** (1986). Hormonal control of sex ratio. *Journal of Theoretical Biology* **118**, 427–441.
- James, W. H.** (1996). Evidence that mammalian sex ratios at birth are partially controlled by parental hormone levels at the time of conception. *Journal of Theoretical Biology* **180**, 271–286.
- Kaňková, Š., Šulc, J., Nouzová, K., Fajfrlík, K., Frynta, D. and Flegr, J.** (2007). Women infected with parasite *Toxoplasma* have more sons. *Naturwissenschaften* **94**, 122–127.

- Kodym, P., Machala, L., Rohacova, H., Sirocka, B. and Maly, M.** (2007). Evaluation of a commercial IgE ELISA in comparison with IgA and IgM ELISAs, IgG avidity assay and complement fixation for the diagnosis of acute toxoplasmosis. *Clinical Microbiology and Infection* **13**, 40–47.
- Lindová, J., Novotná, M., Havlíček, J., Jozífková, E., Skallová, A., Kolbeková, P., Hodný, Z., Kodym, P. and Flegr, J.** (2006). Gender differences in behavioural changes induced by latent toxoplasmosis. *International Journal for Parasitology* **36**, 1485–1492.
- Lutchmaya, S., Baron-Cohen, S., Raggatt, P., Knickmeyer, R. and Manning, J. T.** (2004). 2nd to 4th digit ratios, fetal testosterone and estradiol. *Early Human Development* **77**, 23–28.
- Neave, N., Laing, S., Fink, B. and Manning, J. T.** (2003). Second to fourth digit ratio, testosterone and perceived male dominance. *Proceedings of the Royal Society of London, B* **270**, 2167–2172.
- Okten, A., Kalyoncu, M. and Yaris, N.** (2002). The ratio of second- and fourth-digit lengths and congenital adrenal hyperplasia due to 21-hydroxylase deficiency. *Early Human Development* **70**, 47–54.
- Pokorný, J., Fruhbauer, Z., Polednakova, S., Sykora, J., Zastera, M. and Fialova, D.** (1989). Stanovení antitoxoplasmických protilátek IgG netodou ELISA (Assessment of antitoxoplasmatic IgG antibodies with the ELISA method). *Ceskoslovenska Epidemiologie* **38**, 355–361.
- Roberts, C. W., Walker, W. and Alexander, J.** (2001). Sex-associated hormones and immunity to protozoan parasites. *Clinical Microbiology Reviews* **14**, 476–488.
- Schuster, J. P. and Schaub, G. A.** (2001). Experimental Chagas disease: the influence of sex and psychoneuroimmunological factors. *Parasitology Research* **87**, 994–1000.
- Szczyпка, M. S., Zhou, Q. Y. and Palmiter, R. D.** (1998). Dopamine-stimulated sexual behavior is testosterone dependent mice. *Behavioral Neuroscience* **112**, 1229–1235.
- Tenter, A. M., Heckeroth, A. R. and Weiss, L. M.** (2000). *Toxoplasma gondii*: from animals to humans. *International Journal for Parasitology* **30**, 1217–1258.
- Warren, J. and Sabin, A. B.** (1942). The complement fixation reaction in toxoplasmic infection. *Proceedings of the Society for Experimental Biology and Medicine* **51**, 11–16.
- Webster, J. P.** (2001). Rats, cats, people and parasites: the impact of latent toxoplasmosis on behaviour. *Microbes and Infection* **3**, 1037–1045.