



Photo: André Künzelmann, UFZ

Land use control of nitrate export behavior across catchments


Andreas Musolff¹, Christian Schmidt¹, Benny Selle² and Jan H. Fleckenstein¹

¹ Dep. Hydrogeology, UFZ – Helmholtz-Centre for Environmental Research, Leipzig, Germany

² Institute of Earth and Environmental Science, University of Potsdam, Germany


Introduction – Nutrient exports

- EU Water Framework Directive demands good status of water bodies
 - Elevated nutrients from agriculture lead to eutrophication
 - Catchments as „natural“ management units
- Concentration and load dynamics
- But: complexity of catchment structure and the multitude of the processes involved
- Top-down, data-driven analysis of integrated catchment responses




EEA

Percent of classified groundwater bodies with poor chemical status



Introduction – Nitrate export behavior

- Basu et al. (2010), Thompson et al. (2011): Temporal invariance (chemostatic export regime) of nitrate from managed catchments



Basu et al. (2010), Annual flow weighted concentrations, Baltic Sea Drainage Basin

- Export controlled by discharge
- High availability/ large nitrate store in catchment
- Trajectory in time to develop to this state


Objectives

- Transfer approach to a group of data-rich adjacent catchments with different degrees of agricultural management
→ Good databasis for C/ Q, land use, geology, climatic conditions

Hypothesis: Nitrate export regime is predictable from catchment characteristics and foremost driven by the share of agricultural land use within the catchments



Study area

- Seven sub-catchments within River Bode catchment, two adjacent catchments
- gradients in land use, geology, and climatic conditions
- 16 years time series of NO_3 concentration and discharge (n=74-159)




Methodology – Metrics of export regime

- CV_C/CV_Q → C variance relative to Q variance
- b in $\log C = \log Q * b + d$ → direction of relation between C and Q



$$CV_Q = 0.99$$
$$CV_C = 0.71$$
$$CV_C/CV_Q = 0.72$$

$$b = 0.73$$



Methodology – Partial least squares regression analysis

- Relevance of seven catchment characteristics as predictors for median NO₃ concentrations and metrics of export regime
- But: strong collinearity of catchment characteristics




Discharge per unit area						
0.97**	Runoff coefficient					
-0.85**	-0.72*	Share of drained ara. land				
0.80**	0.79*		topo. gradient			
-0.94**	-0.97**		-0.80**	Share of ara. land		
-0.75*	-0.83**			0.87**	BFI	
-0.86**	-0.91**		-0.75*	0.96**	0.83**	Available water content

PLSR: finding covariance structure in predicting and responding variables

- Can handle collinearity
- Can handle large number of predicting variables
- Interpretation using VIP (variable influence on projection) ranking and regression coefficients



Results – Concentrations & metrics of nitrate export regime

- Export regimes from chemostatic to chemodynamic with positive C-Q relation
- R^2 of logC-logQ relation increases with increasing slope b





Results – PLSR analysis

- Median concentrations (R^2 0.97) and slope b (R^2 0.72) can be well reproduced by catchment characteristics
- CV_C/CV_Q cannot be reproduced (R^2 0.35)





Discussion

- Median nitrate concentrations are driven by agricultural land use
 - Export regimes driven by arable land being artificially drained
- Spatial correlation of nitrate source zone and discharge producing zone is the dominant control of export regime




Conclusions

- Export regime is predictable from catchment characteristics
- Starting hypothesis? → High nitrate inputs and high store does not necessarily lead to chemostatic export regimes
- Trajectory in time from pristine conditions to managed agricultural catchments may be replicated in space
- Endpoint in heavily managed catchments with high degree in hydrological/ land use homogenization



Outlook

- Approach transferable to other catchments and solutes



Musolff et al. (in prep.)