

# Haemosporidian parasites of a European passerine wintering in South Asia: diversity, mixed infections and effect on host condition

P. Synek · T. Albrecht · M. Vinkler · J. Schnitzer ·  
J. Votýpka · P. Munclinger

Received: 18 January 2013 / Accepted: 25 January 2013 / Published online: 6 February 2013  
© Springer-Verlag Berlin Heidelberg 2013

**Abstract** We studied haemosporidian parasites in the scarlet rosefinch *Carpodacus erythrinus* in a small isolated semicolony during an eight-year period using molecular methods of parasite detection. The scarlet rosefinch is an interesting model of parasite host species. It winters in South Asia which represents a rare exception among European passerines. Males express yellow to red carotenoid-based plumage ornament which is a good predictor of male reproductive success. In 240 blood samples originating from 199 adult individuals, the total parasite prevalence reached 60 %. Prevalence varied among years from 36 to 81 % in *Haemoproteus*, 8 to 22 % in *Plasmodium*, and 0 to 14 % in *Leucocytozoon*. Twenty parasite lineages were detected (*Haemoproteus*: 5 lineages, *Plasmodium*: 10 lineages, and *Leucocytozoon*: 5 lineages). Among them, the *Haemoproteus* ROFI2 lineage, which is a host-specific parasite lineage of the scarlet rosefinch, was the most frequently found.

Parasite lineages showed varying degree of lineage specificity. While *Haemoproteus* lineages detected in the scarlet rosefinch have relatively narrow host breadth restricted mainly to Fringillidae family, *Leucocytozoon* and *Plasmodium* lineages generally showed wider host range. The presence of some parasite lineages hitherto detected in sedentary European passerines (SISKIN1, CCF3, BT2) or in *Culicoides* biting midges at the same locality (ROFI1) suggest local transmission. On the contrary, lineages LK05 and FANTAIL1 that were previously reported exclusively from Asian hosts imply parasite transmission at the scarlet rosefinch wintering sites in South Asia. Mixed infections were found in 17 % of infected samples and comprised mainly the most frequent lineages. The pattern of concomitant infections seemed to be rather random and matched expected levels based on lineage frequencies. Between-year comparisons revealed that in a majority of the repeatedly captured individual hosts the infection status remained unchanged (individuals stayed uninfected or possessed the same parasite lineages). However, 16 gains and 8 losses of lineages were also reported. We have not found any effect of haemosporidians on male carotenoid ornament expression or host body mass.

P. Synek · T. Albrecht · M. Vinkler · J. Schnitzer ·  
P. Munclinger (✉)  
Department of Zoology, Charles University in Prague,  
Faculty of Science, Viničná 7,  
128 44 Prague 2, Czech Republic  
e-mail: munclinger@natur.cuni.cz

P. Synek  
e-mail: synek85@gmail.com

T. Albrecht  
e-mail: albrecht@ivb.cz

M. Vinkler  
e-mail: vinkler1@natur.cuni.cz

J. Schnitzer  
e-mail: jan.schnitzer@centrum.cz

J. Votýpka  
Department of Parasitology, Charles University in Prague,  
Faculty of Science, Viničná 7,  
128 44 Prague 2, Czech Republic  
e-mail: vapid@natur.cuni.cz

## Introduction

Haemosporidians, frequent blood parasites of birds transmitted by blood-sucking Dipterian vectors, are diverse group comprising more than 200 described species classified into several genera: *Plasmodium*, *Haemoproteus* and *Leucocytozoon*, which can cause bird disorders (Bennett et al. 1993). Molecular methods of parasite detection (Bensch et al. 2000; Waldenström et al. 2002; Richard et al. 2002; Fallon et al. 2003; Hellgren et al. 2004) started exponential grow of published studies covering various aspects of

haemosporidian diversity, biology, host specificity, and effect on host fitness. Despite this extensive effort, our current view on many aspects of the host–parasite relationships (including parasite lineage specificity and the effect of parasites on host condition) has remained unclear.

Haemosporidian cytochrome *b* lineages which were shown to represent distinct evolutionary entities (Bensch et al. 2004, Valkiūnas et al. 2008a, Valkiūnas et al. 2008b) differ considerably in avian-host specificity. While *Plasmodium* lineages often develop in a large range of phylogenetically unrelated hosts (Waldenström et al. 2002; Križanauskienė et al. 2006; Hellgren et al. 2007; Beadell et al. 2009) and are able to spread between continents via frequent shifts from resident to migratory hosts (Hellgren et al. 2007), *Haemoproteus* lineages tend to show narrower host specificity (Bensch et al. 2009). Moreover, large variability in host specificity exists even among members of each genus. *Plasmodium* lineage SGS1 (assigned to *Plasmodium relictum* morphospecies) infecting almost 50 host species classified into several bird orders and some *Haemoproteus* or *Plasmodium* lineages found in single host species (Bensch et al. 2009) can serve as examples of the extreme limits in parasite–host specialization variability. Typically, Haemosporidian lineage spectra found in a particular host species consist of only few abundant parasite lineages and an array of infrequent lineages. These rare lineages may sometimes represent accidental transmissions into new hosts that cannot result in full development of the parasite and thus form a dead end in the parasite cycle (Valkiūnas et al. 2009a, Valkiūnas 2011). Since the rare infections are frequently transmitted to migratory birds at their wintering grounds where the regular local resident hosts may be known, lineage identity can provide interesting insights into the migratory host biology, especially in species in which the migration pathways are poorly documented.

Experiments performed under controlled laboratory conditions using mainly *Plasmodium* lineages repeatedly showed important pathogenic impact of haemosporidians on their avian hosts (Valkiūnas et al. 2006; Zehntindjiev et al. 2008; Palinauskas et al. 2011). However, the situation in nature seems to be more complex. While some studies demonstrated a significant effect of these haematobious parasites on bird fitness and/or condition (e.g. Seutin 1994; Merila and Sheldon 1999; Merino et al. 2000, Hōrak et al. 2001; Marzal et al. 2005, Westerdahl et al. 2005; Kilpatrick et al. 2006, Knowles et al. 2010), other studies failed to find any effect of this kind (e.g. Dale et al. 1996; Gonzalez et al. 1999; Votýpka et al. 2003; Deviche et al. 2005; Gibb et al. 2005; Bensch et al. 2007). This incongruity probably stems from the following reasons: (1) wild birds usually exhibit low levels of parasitemia suggesting chronic phase of infection which can only slightly affect the host (Valkiūnas 2005, Asghar et al. 2011), (2) while the parasite transmission, which is followed by acute phase of disease, frequently

occurs at wintering grounds, host fitness or condition are measured mostly during the breeding season (Bensch et al. 2007), (3) parasites may be absent in peripheral blood in certain stages of their development and, thus, their detection using blood samples may be problematic (Zehntindjiev et al. 2008), (4) mixed infections which complicate the analyses are frequent (Marzal et al. 2008), and (5) studies may be biased by lower probability of sampling infected then uninfected birds (Valkiūnas 2005).

In our study, we have focused on investigation of haemosporidian parasites in a small community of scarlet rosefinch (*Carpodacus erythrinus*) breeding in the Czech Republic. In eight successive years, we trapped and sampled nearly all individuals at the breeding ground. Scarlet rosefinch is a small, semi-colonial, sexually dimorphic cardueline finch with delayed plumage maturation in males (Stjernberg 1979). Species breeding range spans from central Europe to Kamchatka and covers large area of the Palearctic temperate zone. In the Czech Republic, the scarlet rosefinch breeds at the westernmost limit of its distribution which has been established by recent expansion of the species breeding range (Cramp et al. 1994). The scarlet rosefinch represents an interesting model of a haemosporidian host species for the following two reasons: (1) it is sexually dichromatic, with yellow to red carotenoid-based plumage of males signaling condition and male reproductive success (Albrecht et al. 2007, Vinkler et al. 2012). Hence, the male quality can be easily assessed, and the influence of parasites on male sexual traits can be studied. (2) Avian malaria research is strongly biased towards European–African and American migratory system. Scarlet rosefinch wintering grounds are located in South Asia which represents a rare exception among passerines breeding in Europe. Among birds breeding in Europe and wintering in South Asia, the paddyfield warbler *Acrocephalus agricola* has been the only species studied extensively for avian malaria using molecular methods (Zehntindjiev et al. 2009). Haemosporidian parasites in the scarlet rosefinch have been so far studied only marginally within the framework of comparative studies dealing with limited number of rosefinch individuals (Beadell et al. 2006, Križanauskienė et al. 2006; Hellgren et al. 2007, Pérez-Tris et al. 2007). Four haemosporidian lineages have been hitherto detected in the scarlet rosefinch: *Haemoproteus* ROFI1 and ROFI2, and *Plasmodium* BT7 a SGS1. In our study population, we previously described the impact of *Haemoproteus* infection (detected by microscopy) on individual haematological state (Vinkler et al. 2010). In the present study, we used molecular methods to detect and precisely identify haemosporidian lineages in the same breeding population, estimate parasite prevalence and its temporal variation, reveal mixed infections, and examine the impact of infections on host body mass and male ornamentation.

## Material and methods

### Fieldwork

As the locality and field procedures were described in detail elsewhere (Albrecht 2004; Albrecht et al. 2007 and 2009; Vinkler et al. 2012), we mention here both only briefly. Rosefinches were studied from May to July during the years 2001–2008 at an isolated patch of wet shrubby meadow situated in the Vltava river valley (48°49' N; 13°56' E) in the Bohemia Forest National Park in Doudlebia, Czech Republic. Since the vegetation suitable for rosefinch nesting was searched systematically several times at the locality every year during the egg-laying period, almost all nests have been detected. Adults were mist-netted upon arrival to the locality or during the nest provisioning phase. Caught birds were transported into a field laboratory where basic measurements were taken (weight, tarsus length) and small amount of blood was obtained via brachial venipuncture and stored in ethanol. Breast patch colour (as the most important male ornamentation trait in this species) of males in their third year or older was measured using digital photography and subsequent computer analyses (all described in Albrecht et al. 2009). Shortly, photographs were taken under standard conditions and colour and grey charts were used to further standardize the measurements of colours. Hue, saturation and lightness of carotenoid ornament were measured (HSB colour space) using ADOBE PHOTOSHOP software (Adobe Systems Inc., San Jose, California). We also collected 17 individual blood samples of 13 other bird species trapped accidentally at the study plot (Appendix 1).

### Parasite detection

All blood samples were dried in the laboratory and DNA was extracted using DNeasy® Tissue Kit (Qiagen). Presence and quality of the extracted host (Rosefinch) DNA was inspected by spectrophotometer NanoDrop® ND-1000 (Isogen Life Science) and by control PCRs using mitochondrial DNA (control region and ND2) primers (Pavlova et al. 2005) followed by agarose gel electrophoresis. Parasites were detected following the nested PCR protocol described in Hellgren et al. 2004 (see also Synek et al. 2013 for details), which enables to distinguish *Plasmodium* or *Haemoproteus* infections from *Leucocytozoon* ones using genera-specific nested primers. Positive samples were sequenced using primers HaemF (designed for *Plasmodium* or *Haemoproteus*) or HaemFL (designed for *Leucocytozoon*) (Hellgren et al. 2004). Parasite lineages were identified and classified according to MalAvi database (Bensch et al. 2009). All new haplotypes differing by one or more

substitutions from the sequences deposited in GenBank were sequenced also from the 3'-end with primers HaemR2 and HaemR2L designed for *Plasmodium* or *Haemoproteus* and *Leucocytozoon* respectively. Since the two previously described haemosporidian lineages found in the scarlet rosefinch were named ROFI1 and ROFI2 (using contraction of the word rosefinch), we followed this habit and started the names of the new lineages with ROFI followed by numbers from 3 to 7 as recommended by Pérez-Tris et al. (2007). Sequences of the new lineages were deposited in GenBank (Accession numbers from JX556907 to JX556911).

The basic protocol used (Hellgren et al. 2004) enables simple detection of mixed infection of parasite lineages belonging to different genera (infection by a *Leucocytozoon* lineage and concurrently by *Plasmodium* or *Haemoproteus* lineage). However, mixed infections of *Haemoproteus* and *Plasmodium* lineages or mixed infections of single genus lineages require additional treatment (Loiseau et al. 2008; Marzal et al. 2008). Two strategies were adopted to distinguish these infections: (1) we checked carefully chromatograms for double peaks and (2) we designed lineage-specific primers (Table 1). PCR product of the initial PCR from standard protocol (Hellgren et al. 2004) was used as a template for nested amplifications with lineage-specific primers. Thermal conditions were the same as for the nested PCR in the standard protocol (Hellgren et al. 2004). In each examined sample, presence of the PCR product was verified using an agarose gel. The lineage-specific-primers strategy allowed us to identify mixed infections of *Plasmodium* and *Haemoproteus* lineages and also to detect mixed infections of the most common *Haemoproteus* lineage ROFI2 with other *Haemoproteus* lineages.\

### Statistic analysis and other computations

We used generalized linear models (GLM) with identity link function (assuming normal distribution of dependent variables) for statistical evaluation of host–parasite interactions. Full models with reasonable two-way interactions involving the sex of the birds (where both sexes were treated) were fitted. Simplification of full models started with removing interactions and then the main effect (if they were not part of significant interaction). Hence, model terms were removed gradually, one by one, and models with and without the term of interest were compared using likelihood-ratio tests and *F* statistics (Crawley 2007). Presented are minimal adequate models (MAM) that means model with all members significant ( $\alpha < 0.05$ ). Each individual bird was used only once for the analysis (for those who were captured repeatedly, we used the infection status and other data from the first year when an individual was investigated). Analyses were made in R 2.14.2 software (Mathsoft 2011).

**Table 1** Lineage-specific primers

| Primer name | Sequence (from 5' to 3')    | Product size (bp) | Parasite lineage(s) amplified                            |
|-------------|-----------------------------|-------------------|--|
| SelPlasF    | CATGCAACWGGTGCWTCATT        | 271               | <i>Plasmodium</i> lineages                               |
| SelPlasR    | TTTTTAAGGTTGGGTCACCTACAAG   |                   |  |
| SelHaemF    | ATTGTTACYGCTTTYATGGGTTA     | 150               | <i>Haemoproteus</i> lineages                             |
| SelHaemR    | TCTTTTTAAAGTTGGATCACTWATAGT |                   |  |
| SelH62F     | ATATGCATGCTACTGGAGCTA       | 477               | <i>Haemoproteus</i> ROFI2 Lineage                        |
| SelH62R     | AATAAACTTTGTGCTAAAAATATA    |                   |  |
| SelH392F    | TGCTACCGGTGCTACATTTG        | 471               | <i>Haemoproteus</i> lineages CCF3, SISKIN1, ROFI1, ROFI3 |
| SelH392R    | AATAAACTTTGTGCTAGAAATAGG    |                   |  |

## Results

Two hundred forty blood samples originating from 199 adult scarlet rosefinch individuals (some birds were captured repeatedly in different years) were analysed. We detected at least one haemosporidian lineage in 145 (60 %) samples. The total number of infections was even higher due to mixed infections (Table 2). Sequencing revealed 20 parasite lineages (Table 2) belonging to three haemosporidian genera: *Haemoproteus* (5 lineages),

*Plasmodium* (10 lineages) and *Leucocytozoon* (5 lineages). Five of these lineages were newly identified and these differed by one to eight substitutions from the lineages described previously (Table 3). Individual lineages varied considerably in their prevalence: from dominant ROFI2 which was found in almost half of the samples analysed (112 positive samples, 47 %) to eight rare lineages in which each was detected in a single sample. Apart from ROFI2, only two lineages (SGS1 and BT2) exceeded the level of 5 % prevalence (both lineages 5.4 %). Hence, the pattern of

**Table 2** Haemosporidian lineages found in the scarlet rosefinch. Number of infected individuals is given for particular lineage and year

| Lineage/year (no. of samples) | 2001 (28) | 2002 (24) | 2003 (21) | 2004 (22) | 2005 (37) | 2006 (34) | 2007 (37) | 2008 (37) | Total (240) |
|-------------------------------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-------------|
| <i>Haemoproteus</i>           |           |           |           |           |           |           |           |           |             |
| ROFI2                         | 10        | 11        | 15        | 15        | 14        | 13        | 22        | 12        | 112         |
| CCF3                          |           | 1         | 1         | 1         | 1         |           |           |           | 4           |
| SISKIN1                       |           |           |           | 1         |           |           |           |           | 1           |
| ROFI1                         |           |           | 1         |           |           |           |           | 2         | 3           |
| ROFI3                         |           |           |           |           | 1         |           | 1         |           | 2           |
| <i>Plasmodium</i>             |           |           |           |           |           |           |           |           |             |
| SGS1                          | 2         | 1         | 1         | 2         | 1         | 2         | 1         | 3         | 13          |
| PADOM02                       |           |           |           |           | 2         | 1         | 2         |           | 5           |
| WW3                           | 1         | 1         |           |           | 1         | 1         |           | 1         | 5           |
| ROFI4                         |           | 1         |           |           |           |           |           |           | 1           |
| TURDUS1                       |           |           |           |           |           |           |           | 1         | 1           |
| ROFI5                         |           |           |           |           |           |           |           | 1         | 1           |
| BT6                           |           |           |           |           |           |           |           | 2         | 2           |
| FANTAIL01                     |           | 1         | 1         |           |           |           |           |           | 2           |
| LK05                          |           |           |           |           |           | 1         |           |           | 1           |
| BT8                           |           |           |           |           | 1         |           |           |           | 1           |
| <i>Leucocytozoon</i>          |           |           |           |           |           |           |           |           |             |
| BT2                           | 2         | 1         | 1         | 1         |           | 3         | 4         | 1         | 13          |
| BT5                           |           | 1         |           |           |           |           |           |           | 1           |
| EMSP005                       |           |           |           |           |           | 1         |           |           | 1           |
| ROFI6                         |           |           |           | 1         |           |           |           | 1         | 2           |
| ROFI7                         |           |           |           |           |           |           | 1         | 1         | 2           |
| Total                         | 15        | 18        | 20        | 21        | 21        | 22        | 31        | 25        | 173         |

**Table 3** New lineages

| New lineage                | Closest formerly described lineage | Closest lineage host(s)                     | Difference between lineages |
|----------------------------|------------------------------------|---|-----------------------------|
| <i>Haemoproteus</i> ROFI3  | DENPEN02                           | <i>Dendroica pensylvanica</i> <sup>a</sup>  | 3 substitutions (0.6 %)     |
| <i>Plasmodium</i> ROFI4    | MOALB02                            | <i>Motacilla alba</i> <sup>b</sup>          | 1 substitution (0.2 %)      |
|                            |                                    | <i>Acrocephalus orientalis</i> <sup>b</sup> |                             |
| <i>Plasmodium</i> ROFI5    | MYITYR01                           | <i>Myiarchus tyrannulus</i> <sup>b</sup>    | 6 substitution (1.2 %)      |
| <i>Leucocytozoon</i> ROFI6 | EMSPO04                            | <i>Emberiza spodocephala</i> <sup>c</sup>   | 3 substitutions (0.6 %)     |
|                            | EMSPO05                            |   |                             |
| <i>Leucocytozoon</i> ROFI7 | SILUT01                            | <i>Sicalis luteola</i> <sup>d</sup>         | 8 substitutions (1.7 %)     |

<sup>a</sup> Ricklefs and Fallon 2002; Outlaw and Ricklefs 2009

<sup>b</sup> Beadell et al. 2006

<sup>c</sup> Palinauskas et al. unpublished

<sup>d</sup> Merino et al. 2008


lineage prevalence is markedly right-skewed. Parasite prevalence varied among years from 36 to 81 % in *Haemoproteus*, 8 to 22 % in *Plasmodium*, and 0 to 14 % in *Leucocytozoon* (Fig. 1).

While single-parasite lineage was found in 120 blood samples (83 % of positive samples), mixed infections were detected in 25 samples (17 %). Two lineages were found in most samples showing mixed infections (22 samples); however, three different lineages (one *Haemoproteus*, one *Plasmodium*, and one *Leucocytozoon* lineage) were found in three samples corresponding to three different birds (two females and one male). Mixed infection usually comprised the most frequent lineages (ROFI2 96 % of mixed infections, SGS1 24 %, BT2 24 %). Co-occurrence of lineages in mixed infections matched closely expected values calculated from lineage prevalence (Table 4). Expected total number of concomitant infections (cases when particular lineage shares its host with other lineage) did not differ significantly from observed value (expected 29, observed 28).

Thirty birds were trapped repeatedly (41 retraps) in consecutive years. The interval between repeated trappings of a

particular individual spanned from 1 to 3 years. Four birds which occurred at locality in two different years remained uninfected. Three of them were re-trapped after two years (which suggests they bred one year outside our study plot). It means they were exposed to potential transmission for at least three breeding and wintering seasons. Apart from these uninfected birds, between-year comparisons revealed that in 20 infected individuals the infection status remained unchanged (they harboured the same lineages). However, we also detected 16 gains and 8 losses of lineages in host bloodstream. Losses involved lineages of all three genera (five lineage losses of *Haemoproteus*; one *Plasmodium*, two *Leucocytozoon*). Losses and gains also comprised lineage replacements: one ROFI2 loss was accompanied by gain of BT2, once FANTAIL01 was replaced by ROFI2, and one individual possessing originally ROFI2, BT2, and SGS1 was negative for ROFI2, BT2 and positive for SGS1 and ROFI1 in the next year.

Close inspection of the scarlet rosefinch lineages in the MalAvi database (Bensch et al. 2009) showed varying degree of lineage specificity (Appendix 2). Apart from five

**Fig. 1** Temporal changes of parasite prevalence

**Table 4** Co-occurrence of parasite lineages. Number of mixed infections is given for particular lineage combination. Expected numbers based on lineage prevalence are given in brackets

|       | BT2                | BT5                | ROFI6 | ROFI7              | SGS1               | PADOM02 | WW3  | ROFI4              | TURDUS1            | BT6  | LK5  | CCF3 | ROFI3 |
|-------|--------------------|--------------------|-------|--------------------|--------------------|---------|------|--------------------|--------------------|------|------|------|-------|
| ROFI2 | 6 <sup>a</sup> (6) | 1 <sup>b</sup> (0) | 2(0)  | 2 <sup>c</sup> (1) | 5 <sup>a</sup> (6) | 1(2)    | 1(2) | 1 <sup>b</sup> (0) | 1 <sup>c</sup> (0) | 2(1) | 1(0) | 2(2) | 2(1)  |
| ROFI1 |                    |                    |       |                    | 1(0)               |         |      |                    |                    |      |      |      |       |

<sup>a</sup> Found also in mixed infection of ROFI2, BT2, and SGS1

<sup>b</sup> Found also in mixed infection of ROFI2, BT5, and ROFI4

<sup>c</sup> Found also in mixed infection of ROFI2, ROFI7, and TURDUS1

infrequent novel ROFI lineages, the most frequent lineage ROFI2 has not been reported in any other avian host species. While *Haemoproteus* lineages detected in our study were found exclusively in European passerines of the Fringillidae family (with the exception of SISKIN1 which was found in the house finch *Carpodacus mexicanus* and also the house sparrow *Passer domesticus* in the North America), *Plasmodium* and *Leucocytozoon* lineages show lower host specificity, including even non-passerine birds. They also often exhibit larger geographic distribution. However, *Plasmodium* lineage FANTAIL01 has been hitherto detected only in Australia and Southeast Asia, and *Leucocytozoon* EMSPO05 was known only from black-faced bunting *Emberiza spodocephala* from Russia.

Apart from lineages found in the scarlet rosefinch we detected in this study also six haemosporidian lineages in the other bird species trapped at the study plot (Appendix 1). Among them the *Leucocytozoon* BT2 lineage detected in three bird species was the only lineage shared with the scarlet rosefinch.

We found no association between presence of haemosporidian infections and the expression of condition dependent carotenoid-based colouration of males. The GLM model involving parasite occurrence (*Plasmodium*, *Leucocytozoon*, *Haemoproteus* and the total number of haemosporidian lines found in an individual) as explanatory variables for feather hue was not significant ( $F_{4,93}=0.92$ ,  $P=0.45$ ), the same was true for feather lightness ( $F_{4,93}=0.86$ ,  $P=0.49$ ) and saturation ( $F_{4,93}=0.68$ ,  $P=0.61$ ). In all cases the simplification suggested the null models as the best models.

In both males and females we furthermore evaluated potential associations between body mass (expressed as body weight/tarsus length) and parasite infection. The model involved haemosporidian genera and the total number of lineages, sex and sex interactions with haemosporidian occurrence and lineage numbers as an explanatory variable for body mass. The MAM only included sex as predictor for body mass ( $F_{1,172}=16.5$ ,  $P<0.001$ ), suggesting males were lighter for their tarsus length. However, no effect of haemosporidian infections on body mass was detectable (all  $P>0.15$ ).

## Discussion

We detected relatively high prevalence of haemosporidian parasites in the scarlet rosefinch, if compared to other published studies. However, since we studied only one locality our data may not reflect prevalence for the whole large species range. Prevalence differences have been detected among different populations of a single host species (Bentz et al. 2006, Ferrer et al. 2012) or within one population according to landscape features (Wood et al. 2007). Varying host exposure to insect vectors, which can be hardly assessed in migratory birds that can be infected at their wintering grounds, is probably the most important force determining the level of parasite prevalence (Yohannes et al. 2009). In our study, parasite prevalence varied considerably between years. Unfortunately the timescale was too short to reveal regular pattern of oscillations such were shown in the great reed warbler *Acrocephalus arundinaceus* (Bensch et al. 2007).

Since the migratory birds can be infected at breeding, wintering, and also migratory sites, they are supposed to host larger diversity of parasite lineages than the sedentary birds (Smith et al. 2004). This pattern has been demonstrated repeatedly in various species (e.g. Waldenström et al. 2002; Križanauskienė et al. 2006; Bensch et al. 2007; Hellgren et al. 2007; Ortego et al. 2008, Pérez-Tris et al. 2007). Relatively high number of parasite lineages (four *Haemoproteus* and five *Plasmodium* lineages) has been also detected in the paddyfield warbler (Zehntindjiev et al. 2009), which represents similar (European–South Asian) migration system as the scarlet rosefinch. The number of haemosporidian lineages detected in the scarlet rosefinch supports the view of larger parasite diversity in migrant species and contrasts sharply with only 5 lineages detected in the closely related sedentary house finch (*C. mexicanus*) (Kimura et al. 2006). However, it should be noted that most of the scarlet rosefinch lineages were extremely rare. With the exception of SISKIN1 lineage which was detected also in house sparrow (Marzal et al. 2011) all *Haemoproteus* lineages detected in the scarlet rosefinch have relatively narrow host breadth restricted to Fringillidae family. *Leucocytozoon* and *Plasmodium* lineages generally showed wider host range,

in some *Plasmodium* lineages even also comprising non-passerine species.

Tight co-evolution of parasite lineages and hosts that probably prevents parasites to infect sympatric closely related hosts has been reported repeatedly for avian malaria (Reullier et al. 2006, Pérez-Tris et al. 2007). However, diversity of avian blood parasites was shown not to follow the classical trade-off-hypothesis suggesting that a generalist parasite infecting many hosts should occur at lower prevalence, whilst specialised parasite should rich higher prevalence in the host for which it is specialised (Hellgren et al. 2009). In contrast, Hellgren et al. (2009) showed that both *Haemoproteus* and *Plasmodium* lineages exhibiting the widest host range also occurred at the highest prevalence in a single host species. In concordance with these results, the most frequent *Leucocytozoon* and *Plasmodium* lineages detected in the present study had, indeed, the widest host ranges (*Plasmodium* SGS1 48 host species, *Leucocytozoon* BT2 8 host species). Nevertheless, the most frequent *Haemoproteus* lineage ROFI2, which is the most frequent haemosporidian lineage in the present study and was detected in the scarlet rosefinch also by Križanauskienė et al. (2006) and Bensch et al. (2009), has not been found in any other species, which indicates its strict host specificity.

*Haemoproteus* lineages SISKIN1 and CCF3 have been previously found only in European sedentary species, which suggests their local transmission to the scarlet rosefinch at the breeding site. We may also speculate that BT2 is transmitted during the breeding season, because it was present in other bird species at the study plot including the sedentary willow tit (*Poecile montanus*). Transmission during the scarlet rosefinch breeding can also be suggested for ROFI1 which was found in *Culicoides* biting midges at the same study plot in 2008 (Synek et al. 2013). On the other hand, the most frequent ROFI2 lineage was not found in haematophagous insects trapped at the same locality, which may suggest its transmission at wintering sites (Synek et al. 2013). Lineages that have been hitherto exclusively detected in Asian bird hosts (*Leucocytozoon* EMSPO05 known from *Emberiza spodocephala*, and *Plasmodium* FANTAIL01 detected in *Acridotheres tristis*, *Dendrocygna javanica* and *Rhipidura rufifrons*) correspond well with the assumed scarlet rosefinch wintering sites in South Asia. Since the EMSPO05 and FANTAIL01 have not been detected in any other European bird hosts, it is highly likely that scarlet rosefinches were infected by those lineages at their wintering sites. However, since avian malaria is far less studied in South Asia than in Europe and Africa we cannot exclude the possibility of winter transmission of other lineages that have been hitherto detected only in Europe and Africa.

Mixed infections of haemosporidian lineages have been repeatedly reported in avian hosts (e.g. Marzal et al. 2008; Valkiūnas et al. 2009b; Asghar et al. 2011; Zehtindjiev et al.

2012). Observed number of mixed infections in the present study closely matched the expected values based on prevalence of individual lineages, which may suggest random pattern of concomitant infections. However, it should be noted that the probability of detecting mixed infections strongly depends on the detection methods. For example, while detection based on manual inspection of electropherograms for double peaks suggested that concomitant infections were less frequent than expected in a long-term population study of great reed warblers (Bensch et al. 2007), closer inspection of partially overlapping dataset using specific primers and highly sensitive quantitative PCR showed on contrary significantly higher frequency of mixed infections than expected (Asghar et al. 2011). Even though we used both manual inspection of electropherograms and specific primer approach, we cannot fully exclude the possibility that we underestimated the frequency of mixed infections, especially in cases of low parasitemia in one of the lineages. On the other hand, in such a case the prevalence of individual lineages would be also underestimated in exactly the same way.

Infection status of birds trapped repeatedly in different years remained mostly unchanged. It may be interpreted either as evidence for a long-term survival of parasites in the host's bloodstream or as evidence for variability in individual host sensitivity to particular parasite lineages and consequent recurrent infections (Hasselquist et al. 2007). While the data showing long-term persistence of haemosporidian parasites in the chronic phase of their life cycle in avian hosts (Valkiūnas 2005, Zehtindjiev et al. 2008) support the former explanation, the known association between MHC alleles and resistance to avian malaria (reviewed in Westerdahl 2007) is consistent with the later explanation. On the other hand, relatively high rates of both *Haemoproteus* and *Plasmodium* lineage losses were also recorded in other bird species (*A. arundinaceus*: Bensch et al. 2007, Hasselquist et al. 2007; *Cyanistes caeruleus*: Knowles et al. 2011). In our study, the number of parasite lineage gains (16) twice exceeded lineage losses (8). However, since the *Haemoproteus* (five losses) and *Leucocytozoon* (two losses) lineages life cycles also involve stages in host internal organs, absence of gametocytes in the host bloodstream does not necessarily imply the full host recovery from the infection. Lineage replacements and the only one detected loss of *Plasmodium* lineage can be probably explained by very low parasitemia below the limits of our detection methods. In conclusion, we have not found any strong evidence of complete elimination of parasite lineages and hence our results are compatible with long-time persistence of parasites in bird hosts.

Carotenoid ornaments are generally considered as honest indicators of health (Lozano 1994, Badyaev and Hill 2000,

Vinkler and Albrecht 2010). In the present study, however, we have not found any effect of haemosporidian presence on male coloration and host body mass. In our previous study (Vinkler et al. 2010) based on microscopic examination of blood smears, we have shown that haemosporidian parasites may influence health in the scarlet rosefinch (measured as basophil ratio in peripheral blood). Nonetheless, the present lack of supportive evidence for the haemosporidian influence on body mass is in accordance with number of field and laboratory studies in birds (Sanz et al. 2002; Votýpka et al. 2003, Gibb et al. 2005; Deviche et al. 2005; Palinauskas et al. 2008). It is, thus, possible that the loss of weight due to blood parasites may be only temporary; occurring in acute phase of the infection and birds may recover to their original weight during the chronic phase (Valkiūnas 2005). Studies targeting the effect of haemosporidian parasites on carotenoid-based male coloration are rather scarce and inconsistent. While no effect of haemosporidian parasite presence on male coloration was found in redpoll (*Carduelis flammea*) Seutin (1994), negative correlation between haemosporidian presence and intensity of ornamentation was observed in yellowhammer (*Emberiza citrinella*) (Sundberg 1995). Surprisingly, also positive correlation of parasite presence and male carotenoid ornaments was reported in greenfinch (*Carduelis chloris*) (Merila and Sheldon 1999). Thus, it seems the effect of haemosporidians on carotenoid ornaments can be also rather complex and age dependent, as shown in blue tit (*C. caeruleus*) Hōrak et al. (2001). It should be also noted that the feather ornament of scarlet rosefinches can be influenced mainly during moulting completed during winter. We cannot exclude the possibility that the presence of parasites detected during the breeding season only weakly correlates with the bird infection status in winter. Moreover, acute winter infections that we were unable to detect during their chronic phase in spring may influence the host ornaments. However, this view remains unsupported by the fact that majority of birds trapped repeatedly in different years unchanged their infection status.

**Acknowledgments** The study was supported by Czech Science Foundation (GA ČR) grant No. P506/10/0716. P. Synek received support from the SVV 2013-267 201 project of the Charles University in Prague. We acknowledge the help of our colleagues and friends Jaroslav Jelínek and František Zicha in the field and thank Zdena Csiebreiová for her technical assistance in the laboratory.

**Ethical standards** This study was performed under certificate of competency according to §17 of the Act No. 246/1992 coll. on Protection Animals against Cruelty and comply with the current law of the Czech Republic.

**Conflict of interest** The authors declare that they have no conflict of interest.

## Appendix 1

**Table 5** Lineages found in other species trapped at the scarlet rosefinch locality

| Host species  | No. of individuals | <i>Haemoproteus</i> | <i>Leucocytozoon</i> |
|---|--------------------|---------------------|----------------------|
| Common Treecreeper ( <i>Certhia familiaris</i> )    | 1                  |                     |                      |
| House Martin ( <i>Delichon urbica</i> )             | 1                  |                     | SYBOR8               |
| Chaffinch ( <i>Fringilla coelebs</i> )              | 1                  | CCF1                |                      |
| Red-backed Shrike ( <i>Lanius collurio</i> )        | 2                  | RB1                 | BT2                  |
| Bluethroat ( <i>Luscinia svecica</i> )              | 1                  |                     | BT2                  |
| Willow Tit ( <i>Parus montanus</i> )                | 1                  |                     | BT2                  |
| Common Chiffchaff ( <i>Phylloscopus collybita</i> ) | 1                  |                     |                      |
| Willow Warbler ( <i>Phylloscopus trochilus</i> )    | 1                  |                     |                      |
| Dunnock ( <i>Prunella modularis</i> )               | 2                  | TURDUS2, RB1        |                      |
| Garden Warbler ( <i>Sylvia borin</i> )              | 2                  |                     |                      |
| Common Whitethroat ( <i>Sylvia communis</i> )       | 2                  | CWT3                |                      |
| Lesser Whitethroat ( <i>Sylvia curruca</i> )        | 1                  |                     |                      |
| Blackbird ( <i>Turdus merula</i> )                  | 1                  | TURDUS2             |                      |

## Appendix 2

**Table 6** Lineage specificity. Other hosts of the scarlet rosefinch parasite lineages are listed according to MalAvi database. Six lineages found exclusively in the scarlet rosefinch are not included

| Lineage             | Host species | Geographic region  |
|---------------------|--------------|--|
| <i>Haemoproteus</i> | CCF3         | <i>Fringilla coelebs</i> Europe  |
|                     | SISKIN1      | <i>Carpodacus mexicanus</i> Europe, North America<br><i>Carduelis spinus</i><br><i>Loxia curvirostra</i><br><i>Passer domesticus</i> |
|                     | ROFI1        | <i>Carpodacus erythrinus</i> Europe<br><i>Coccothraustes coccothraustes</i><br><i>Fringilla coelebs</i><br><i>Carduelis chloris</i>  |



**Table 6** (continued)

| Lineage              | Host species  | Geographic region   |
|----------------------|---|---|
| <i>Plasmodium</i>    | SGS1  | 48 host species<br>Europe, Asia, Africa,<br>New Zealand   |
|                      | PADOM02   | <i>Passer domesticus</i><br><i>Anthus hodgsoni</i><br><i>Corvus corone</i><br><i>Luscinia svecica</i><br><i>Motacilla flava</i><br><i>Phasianus colchicus</i><br><i>Emberiza citrinella</i><br><i>Passer montanus</i> |
| WW3                  |   | Europe, North America, Africa   |
|                      |   | <i>Andropadus virens</i>  |
|                      |   | <i>Geothlypis trichas</i>   |
|                      |   | <i>Quelea quelea</i>  |
|                      |   | <i>Phylloscopus trochilus</i>   |
|                      |   | <i>Spermophaga haematina</i>  |
|                      |   | <i>Quelea quelea</i>  |
|                      |   | <i>Phylloscopus trochilus</i>   |
|                      |   | <i>Passer montanus</i>  |
|                      |   | <i>Passer domesticus</i>  |
| TURDUS1              | 18 host species   | Europe, South Asia, Africa  |
|                      |   | <i>Luscinia svecica</i>   |
| BT6                  |   | Europe  |
| FANTAIL01            | <i>Acridotheres tristis</i><br><i>Dendrocygna javanica</i><br><i>Rhipidura rufifrons</i>  | Australia, South Asia   |
|                      |   |   |
|                      |   |   |
| LK05                 | <i>Falco namanni</i>  | Europe  |
| BT8                  | <i>Luscinia svecica</i><br><i>Ploceus</i> 2 species<br><i>Hypothymis azurea</i><br><i>Estrilda amandava</i><br><i>Copsychus</i> 2 species<br><i>Lophura punctuata</i> | Europe, Africa, South Asia  |
|                      |   |   |
|                      |   |   |
|                      |   |   |
|                      |   |   |
|                      |   |   |
| <i>Leucocytozoon</i> | BT2   | Europe, Africa  |
|                      |   | <i>Lanius collurio</i><br><i>Muscicapa striata</i><br><i>Phoenicurus phoenicurus</i>  |
|                      |   |   |

**Table 6** (continued)

| Lineage | Host species                                      | Geographic region |
|---------|---|-------------------|
|         | <i>Phylloscopus trochilus</i>                     |                   |
|         | <i>Saxicola rubetra</i>                           |                   |
|         | <i>Sylvia atricapilla</i>                         |                   |
|         | <i>Sylvia borin</i>                               |                   |
|         | <i>Hippolais icterina</i>                         |                   |
| BT5     | <i>Luscinia svecica</i><br><i>Lanius collurio</i> | Europe            |
| EMSP005 | <i>Emberiza spodocephala</i>                      | Asia              |

## References

- Albrecht T (2004) Edge effect in wetland-arable land boundary determines nesting success of scarlet rosefinches (*Carpodacus erythrinus*) in the Czech Republic. *Auk* 121:361–371
- Albrecht T, Schnitzer J, Kreisinger J, Exnerová A, Bryja J, Munclinger P (2007) Extrapair paternity and the opportunity for sexual selection in long-distant migratory passerines. *Behav Ecol* 18:477–486
- Albrecht T, Vinkler M, Schnitzer J, Poláková R, Munclinger P, Bryja J (2009) Extra-pair fertilizations contribute to selection on secondary male ornamentation in a socially monogamous passerine. *J Evol Biol* 22:2020–2030
- Asghar M, Hasselquist D, Bensch S (2011) Are chronic avian haemosporidian infections costly in wild birds? *J Avian Biol* 42:530–537
- Badyaev AV, Hill GE (2000) Evolution of sexual dichromatism: contribution of carotenoid- versus melanin-based coloration. *Biol J Linn Soc* 69:153–172
- Beadell JS, Covas R, Gebhard C, Ishtiaq F, Melo M, Schmidt BK, Perkins SL, Graves GR, Fleischer RC (2009) Host associations and evolutionary relationships of avian blood parasites from West Africa. *Int J Parasitol* 39:257–266
- Beadell JS, Ishtiaq F, Covas R, Melo M, Warren BH, Atkinson CT, Bensch S, Graves GR, Jhala YV, Peirce MA, Rahmani AR, Fonseca DM, Fleischer RC (2006) Global phylogeographic limits of Hawaii's avian malaria. *P Roy Soc Lond B Bio* 273:2935–2944
- Bensch S, Hellgren O, Pérez-Tris (2009) MalAvi: a public database of malaria parasites and related haemosporidians in avian hosts based on mitochondrial cytochrome b lineages. *Mol Ecol Resour* 9:1353–1358
- Bensch S, Pérez-Tris J, Waldenström J, Hellgren O (2004) Linkage between nuclear and mitochondrial DNA sequences in avian malaria parasites: Multiple cases of cryptic speciation? *Evolution* 58:1617–1621
- Bensch S, Stjernman M, Hasselquist D, Ostman O, Hansson B, Westerdaal H, Pinheiro RT (2000) Host specificity in avian blood parasites: a study of *Plasmodium* and *Haemoproteus* mitochondrial DNA amplified from birds. *P Roy Soc Lond B Bio* 267:1583–1589
- Bensch S, Waldenström J, Jonzen N, Westerdaal H, Hansson B, Sejberg D, Hasselquist D (2007) Temporal dynamics and diversity of avian malaria parasites in a single host species. *J Anim Ecol* 76:112–122

- Bennett GF, Peirce MA, Ashford RW (1993) Avian Haematozoa—mortality and pathogenicity. *J Nat Hist* 27:993–1001
- Bentz S, Rigaud T, Barroca M, Martin-Laurent F, Bru D, Moreau J, Faivre B (2006) Sensitive measure of prevalence and parasitaemia of haemosporidia from European blackbird (*Turdus merula*) populations: value of PCR-RFLP and quantitative PCR. *Parasitology* 133:685–692
- Cramp P, Perrins CM, Brooks DJ (1994) The birds of Western Palearctic. Oxford University Press, Oxford
- Crawley MJ (2007) The R book. Wiley, Chichester
- Dale S, Kruszewicz A, Slagsvold T (1996) Effects of blood parasites on sexual and natural selection in the pied flycatcher. *J Zool* 238:373–393
- Deviche P, McGraw K, Greiner EC (2005) Interspecific differences in hematozoan infection in sonoran desert *Aimophila* sparrows. *J Wildlife Dis* 41:532–541
- Ferrer ES, Garcia-Navas V, Sanz JJ, Ortego J (2012) Molecular characterization of avian malaria parasites in three Mediterranean blue tit (*Cyanistes caeruleus*) populations. *Parasitol Res* 111:2137–2142
- Fallon SM, Ricklefs RE, Swanson BL, Bermingham E (2003) Detecting avian malaria: an improved polymerase chain reaction diagnostic. *J Parasitol* 89:1044–1047
- Gibb CE, Jones J, Girvan MK, Barg JJ, Robertson RJ (2005) Geographic variation in prevalence and parasitemia of *Haemoproteus paruli* in the cerulean warbler (*Dendroica cerulea*). *Can J Zool* 83:626–629
- Gonzalez G, Sorci G, Møller AP, Ninni P, Haussy C, De Lope F (1999) Immunocompetence and condition-dependent sexual advertisement in male house sparrows (*Passer domesticus*). *J Anim Ecol* 68:1225–1234
- Hasselquist D, Östman Ö, Waldenström J, Bensch S (2007) Temporal patterns of occurrence and transmission of the blood parasite *Haemoproteus payevskyi* in the great reed warbler *Acrocephalus arundinaceus*. *J Ornithol* 148:401–409
- Hellgren O, Pérez-Tris J, Bensch S (2009) A jack-of-all-trades and still a master of some: prevalence and host range in avian malaria and related blood parasites. *Ecology* 90:2840–2849
- Hellgren O, Waldenström J, Bensch S (2004) A new PCR assay for simultaneous studies of *Leucocytozoon*, *Plasmodium*, and *Haemoproteus* from avian blood. *J Parasitol* 90:797–802
- Hellgren O, Waldenström J, Pérez-Tris J, Szollosi E, Hasselquist D, Krizanauskienė A, Ottosson U, Bensch S (2007) Detecting shifts of transmission areas in avian blood parasites—a phylogenetic approach. *Mol Ecol* 16:1281–1290
- Hörak P, Ots I, Vellau H, Spottiswoode C, Møller AP (2001) Carotenoid-based plumage coloration reflects hemoparasite infection and local survival in breeding great tits. *Oecologia* 126:166–173
- Kilpatrick AM, LaPointe DA, Atkinson CT, Woodworth BL, Lease JK, Reiter ME, Gross K (2006) Effects of chronic avian malaria (*Plasmodium relictum*) infection on reproductive success of Hawaii Amakihi (*Hemignathus virens*). *Auk* 123:764–774
- Kimura M, Dhondt AA, Lovette IJ (2006) Phylogeographic structuring of *Plasmodium* lineages across the North American range of the house finch (*Carpodacus mexicanus*). *J Parasitol* 92:1043–1049
- Knowles SCL, Palinauskas V, Sheldon BC (2010) Chronic malaria infections increase family inequalities and reduce parental fitness: experimental evidence from a wild bird population. *J Evol Biol* 23:557–569
- Knowles SCL, Wood MJ, Alves R, Wilkin TA, Bensch S, Sheldon BC (2011) Molecular epidemiology of malaria prevalence and parasitaemia in a wild bird population. *Mol Ecol* 20:1062–1076
- Krizanauskienė A, Hellgren O, Kosatec V, Sokolov L, Bensch S, Valkiūnas G (2006) Variation in host specificity between species of avian haemosporidian parasites: evidence from parasite morphology and cytochrome b gene sequences. *J Parasitol* 92:1319–1324
- Loiseau C, Zoorob R, Garnier S, Birard J, Federici P, Julliard R, Sorci G (2008) Antagonistic effects of a Mhc class I allele on malaria-infected house sparrows. *Ecol Lett* 11:258–265
- Lozano GA (1994) Carotenoids, parasites, and sexual selection. *Oikos* 70:309–311
- Marzal A, de Lope F, Navarro C, Møller AP (2005) Malarial parasites decrease reproductive success: an experimental study in a passerine bird. *Oecologia* 142:541–554
- Marzal A, Bensch S, Reviriego M, Balbontin J, de Lope F (2008) Effects of malaria double infection in birds: one plus one is not two. *J Evol Biol* 21:979–987
- Marzal A, Ricklefs RE, Valkiūnas G, Albayrak T, Arriero E, Bonneaud C, Cziráková GA, Ewen J, Hellgren O, Hořáková D, Iezhova TA, Jensen H, Krizanauskienė A, Lima MR, de Lope F, Magnussen E, Martin LB, Møller AP, Palinauskas V, Pap PL, Pérez-Tris J, Sehgal RN, Soler M, Szöllosi E, Westerdaal H, Zetindjiev P, Bensch S (2011) Diversity, loss, and gain of malaria parasites in a globally invasive bird. *PLoS One* 6:e21905
- Merila J, Sheldon BC (1999) Testis size variation in the greenfinch *Carduelis chloris*: relevance for some recent models of sexual selection. *Behav Ecol Sociobiol* 45:115–123
- Merino S, Moreno J, Sanz JJ, Arriero E (2000) Are avian blood parasites pathogenic in the wild? A medication experiment in blue tits (*Parus caeruleus*). *P Roy Soc Lond B Bio* 267:2507–2510
- Merino S, Moreno J, Vásquez RA, Martínez J, Sánchez-Monsálvez I, Estades CF, Ippi S, Sabat P, Rozzi R, McGehee S (2008) Haematozoa in forest birds from southern Chile: latitudinal gradients in prevalence and parasite lineage richness. *Austral Ecol* 33:329–340
- Ortego J, Calabuig G, Cordero PJ, Aparicio JM (2008) Genetic characterization of avian malaria (Protozoa) in the endangered lesser kestrel, *Falco naumanni*. *Parasitol Res* 101:1153–1156
- Outlaw DC, Ricklefs RE (2009) On the phylogenetic relationships of haemosporidian parasites from raptorial birds (Falconiformes and Strigiformes). *J Parasitol* 95:1171–1176
- Palinauskas V, Valkiūnas GN, Bolshakov CV, Bensch S (2008) *Plasmodium relictum* (lineage P-SGS1): effects on experimentally infected passerine birds. *Exp Parasitol* 120:372–380
- Palinauskas V, Valkiūnas G, Bolshakov CV, Bensch S (2011) *Plasmodium relictum* (lineage SGS1) and *Plasmodium ashfordi* (lineage GRW2): The effects of the co-infection on experimentally infected passerine birds. *Exp Parasitol* 127:527–533
- Pavlova A, Zink RM, Rohwer S (2005) Evolutionary history population genetics and gene flow in the common rosefinch (*Carpodacus erythrinus*). *Mol Phylogenet Evol* 36:669–681
- Pérez-Tris J, Hellgren O, Krizanauskienė A, Waldenström J, Secondi J, Bonneaud C, Fjeldsa J, Hasselquist D, Bensch S (2007) Within-host speciation of malaria parasites. *PLoS One* 2:e235
- Reullier J, Perez-Tris J, Bensch S, Secondi J (2006) Diversity, distribution and exchange of blood parasites meeting at an avian moving contact zone. *Mol Ecol* 15:753–763
- Richard FA, Sehgal RNM, Jones HI, Smith TB (2002) A comparative analysis of PCR-based detection methods for avian malaria. *J Parasitol* 88:819–822
- Ricklefs RE, Fallon SM (2002) Diversification and host switching in avian malaria parasites. *Proc R Soc Lond B* 269:885–892
- Sanz JJ, Moreno J, Arriero E, Merino S (2002) Reproductive effort and blood parasites of breeding pied flycatchers: the need to control for interannual variation and initial health state. *Oikos* 96:299–306
- Seutin G (1994) Plumage redness in Redpoll Finches does not reflect hemoparasitic infection. *Oikos* 70:280–286
- Smith RB, Greiner EC, Wolf BO (2004) Migratory movements of Sharp-shinned hawks (*Accipiter striatus*) captured in New Mexico in relation to prevalence intensity and biogeography of avian hematozoa. *Auk* 121:837–846

- Stjernberg T (1979) Breeding biology and population dynamics of the scarlet rosefinch *Carpodacus erythrinus*. Finnish Zoological Pub, Board
- Sundberg J (1995) Parasites, plumage coloration and reproductive success in the yellowhammer. *Emberiza citrinella* OIKOS 74:331–339
- Synek P, Munclinger P, Albrecht T, Votýpka J (2013) Avian haemosporidians in haematophagous insects in the Czech Republic. *Parasitol Res* 112:839–845
- Valkiūnas G (2005) Avian malaria parasites and other haemosporidia. CRC, Boca Raton
- Valkiūnas G (2011) Haemosporidian vector research: marriage of molecular and microscopical approaches is essential. *Mol Ecol* 20:3084–3086
- Valkiūnas G, Zickus T, Shapoval AP, Lezhova TA (2006) Effect of *Haemoproteus belopolnyi* (Haemosporida: Haemoproteidae) on body mass of the blackcap *Sylvia atricapilla*. *J Parasitol* 92:1123–1125
- Valkiūnas G, Iezhova TA, Loiseau C, Chasar A, Smith TB, Sehgal RNM (2008a) New species of haemosporidian parasites (Haemosporida) from African rainforest birds, with remarks on their classification. *Parasitol Res* 103:1213–1228
- Valkiūnas G, Zehtindjiev P, Dimitrov D, Križanauskiene A, Iezhova TA, Bensch S (2008b) Polymerase chain reaction-based identification of *Plasmodium* (Huffia) *elongatum*, with remarks on species identity of haemosporidian lineages deposited in GenBank. *Parasitol Res* 102:1185–1193
- Valkiūnas G, Iezhova TA, Loiseau C, Sehgal RNM (2009a) Nested cytochrome B polymerase chain reaction diagnostics detect sporozoites of haemosporidian parasites in peripheral blood of naturally infected birds. *J Parasitol* 95:1512–1515
- Valkiūnas G, Iezhova TA, Loiseau C, Smith TB, Sehgal RNM (2009b) New malaria parasites of the subgenus *Novyella* in African rainforest birds, with remarks on their high prevalence, classification and diagnostics. *Parasitol Res* 104:1061–1077
- Vinkler M, Albrecht T (2010) Carotenoid maintenance handicap and the physiology of carotenoid-based signalisation of health. *Naturwissenschaften* 97:19–28
- Vinkler M, Schnitzer J, Munclinger P, Albrecht T (2012) Phytohaemagglutinin skin-swelling test in scarlet rosefinch males: low-quality birds respond more strongly. *Anim Behav* 83:17–23
- Vinkler M, Schnitzer J, Munclinger P, Votýpka J, Albrecht T (2010) Haematological health assessment in a passerine with extremely high proportion of basophils in peripheral blood. *J Ornithol* 151:841–849
- Votýpka J, Simek J, Tryjanowski P (2003) Blood parasites, reproduction and sexual selection in the red-backed shrike (*Lanius collurio*). *Ann Zool Fenn* 40:431–439
- Waldenström J, Bensch S, Kiboi S, Hasselquist D, Ottosson U (2002) Cross-species infection of blood parasites between resident and migratory songbirds in Africa. *Mol Ecol* 11:1545–1554
- Westerdahl H (2007) Passerine MHC: genetic variation and disease resistance in the wild. *J Ornithol* 148:S469–S477
- Westerdahl H, Waldenström J, Hansson B, Hasselquist D, von Schantz T, Bensch S (2005) Associations between malaria and MHC genes in a migratory songbird. *P Roy Soc Lond B Biol* 272:1511–1518
- Wood MJ, Cosgrove CL, Wilkin TA, Knowles SCL, Day KP, Sheldon BC (2007) Within population variation in prevalence and lineage distribution of avian malaria in blue tits, *Cyanistes caeruleus*. *Mol Ecol* 16:3263–3273
- Yohannes E, Križanauskiene A, Valcu M, Bensch S, Kempenaers B (2009) Prevalence of malaria and related haemosporidian parasites in two shorebird species with different winter habitat distribution. *J Ornithol* 150:287–291
- Zehtindjiev P, Ilieva M, Westerdahl H, Hansson B, Valkiūnas G, Bensch S (2008) Dynamics of parasitemia of malaria parasites in a naturally and experimentally infected migratory songbird, the great reed warbler *Acrocephalus arundinaceus*. *Exp Parasitol* 119:99–110
- Zehtindjiev P, Ilieva M, Križanauskiene A, Oparina O, Oparin M, Staffan Bensch S (2009) Occurrence of haemosporidian parasites in the paddyfield warbler, *Acrocephalus agricola* (Passeriformes, Sylviidae). *Acta Parasitol* 54:295–300
- Zehtindjiev P, Križanauskiene A, Bensch S, Palinauskas V, Asghar M, Dimitrov D, Scebba S, Valkiūnas G (2012) A new morphologically distinct avian malaria parasite that fails detection by established polymerase chain reaction-based protocols for amplification of the cytochrome B gene. *J Parasitol* 98:657–665