

Doporučená literatura:

- Ackerman, L., Hajná, J., Žák, J., Erban, V., Sláma, J., Polák, L., Kachlík, V., Strnad, L., Trubač, J., 2019. Architecture and composition of ocean floor subducted beneath northern Gondwana during Neoproterozoic to Cambrian: A palinspastic reconstruction based on Ocean Plate Stratigraphy (OPS). *Gondwana Res.* 76, 77-97.
- Ackerman, L., Pašava, J., Žák, J., Žák, K., Kachlík, V., Šebek, O., Trubač, J., Svojtka, M., Veselovský, F., Strnad, L., Santolík, V., 2020. Arc-related black shales as sedimentary archives of sea-level fluctuations and plate tectonics during the late Neoproterozoic: An example from the Bohemian Massif. *Marine and Petroleum Geology*, 104713.
- Hajná, J., Dörr, W., Kachlík, V., Sláma, J., 2018. New constraints from detrital zircon ages on protracted, multiphase transition from the Cadomian accretionary orogen to a passive margin of Gondwana. *Precamb. Res.* 317, 159-173.
- Hajná, J., Žák, J., Kachlík, V., 2011. Structure and stratigraphy of the Tepla-Barrandian Neoproterozoic, Bohemian Massif: A new plate-tectonic reinterpretation. *Gondwana Research* 19, 495-508.
- Hajná, J., Žák, J., Ackerman, L., Svojtka, M., Pašava, J., 2019. A giant late Precambrian chert-bearing olistostrome discovered in the Bohemian Massif: A record of Ocean Plate Stratigraphy (OPS) disrupted by mass-wasting along an outer trench slope. *Gondwana Res.* 74, 173-188.
- Kachlík, V., 1999. Relationship between Moldanubicum, the Kutná Hora Crystalline unit and Bohemium (Central Bohemia, Czech Republic): A result of the polyphase Variscan nappe tectonics. *J. Czech. Geol. Soc.* 44, 201-291.
- Nahodilová, R., Hasalová, P., Štípská, P., Schulmann, K., Závada, P., Míková, J., Kylander-Clark, A., Maierová, P., 2019. Exhumation of subducted continental crust along the arc region. *Gondwana Res.*
- Santolík, V., Ackerman, L., Kachlík, V., Sláma, J., Mészáros, N., 2022. Petrogenesis of low-pressure intra-oceanic arc granitoids: Insights from the late Neoproterozoic Avalonian–Cadomian orogen, Bohemian Massif. *Lithos* 428-429, 106808.
- Sláma, J., Dunkley, D.J., Kachlík, V., Kusiak, M.A., 2008. Transition from island-arc to passive setting on the continental margin of Gondwana: U-Pb zircon dating of Neoproterozoic metaconglomerates from the SE margin of the Tepla-Barrandian Unit, Bohemian Massif. *Tectonophysics* 461, 44-59.
- Soejono, I., Machek, M., Sláma, J., Janoušek, V., Kohút, M., 2019. Cambro–Ordovician anatexis and magmatic recycling at the thinned Gondwana margin: new constraints from the Kouřim Unit, Bohemian Massif. *J. Geol. Soc.*, 2019-2037.
- Žák, J., Verner, K., Janoušek, V., Holub, F., Kachlík, V., Finger, F., Hajná, J., Tomek, F., Vondrovič, L., Trubač, J., 2014. A plate-kinematic model for the assembly of the Bohemian Massif constrained by structural relationship around granitoid pluton, in: Schulmann, K., Martínez Catalán, J.R., Lardeaux, J.M., Janoušek, V., Oggiano, G. (Eds.), *The Variscan Orogeny: Extent, Timescales and the Formation of the European Crust*. Geological Society, London, London.