O APPLICATIONS OF NEXT-GENERATION SEQUENCING

Harnessing ancient genomes to study the history of human adaptation

Stephanie Marciniak¹ and George H. Perry^{1,2,3}

Abstract | The past several years have witnessed an explosion of successful ancient human genome-sequencing projects, with genomic-scale ancient DNA data sets now available for more than 1,100 ancient human and archaic hominin (for example, Neandertal) individuals. Recent 'evolution in action' analyses have started using these data sets to identify and track the spatiotemporal trajectories of genetic variants associated with human adaptations to novel and changing environments, agricultural lifestyles, and introduced or co-evolving pathogens. Together with evidence of adaptive introgression of genetic variants from archaic hominins to humans and emerging ancient genome data sets for domesticated animals and plants, these studies provide novel insights into human evolution and the evolutionary consequences of human behaviour that go well beyond those that can be obtained from modern genomic data or the fossil and archaeological records alone.

Adaptation

A process of phenotypic and corresponding genetic change over time for traits that confer increased reproductive fitness in a given environmental context.

¹Department of Anthropology, Pennsylvania State University, University Park, Pennsylvania 16802, USA. ²Department of Biology, Pennsylvania State University, University Park, Pennsylvania 16802, USA. ³Huck Institutes of the Life Sciences, Pennsylvania State University, University Park, Pennsylvania 16802, USA.

Correspondence to G.H.P. ghp3@psu.edu

doi:<u>10.1038/nrg.2017.65</u> Published online 11 Sep 2017 Reconstructing the history of human biological adaptation has fascinated scientists (and many others) for more than 150 years^{1,2}. How did humans evolve in response to the new environments they encountered while expanding across the globe³? Did prehistoric human populations adapt biologically to cultural changes such as the dietary shifts associated with agriculture⁴? Do our past adaptations have consequences for health and medicine today, especially given the extent to which some modern human environments and diets differ from those of the past?

With the human reference genome as a cornerstone and benefiting from advances in genotyping and sequencing technology, evolutionary population genetic analyses have greatly aided the study of human biological adaptation. That is, on the basis of patterns of genetic variation observed among living people, researchers can identify genome regions that contain putative 'signatures' of past positive natural selection^{5,6}. If the functional consequences of candidate genetic variants can be determined using genotype-phenotype association studies7 or animal or cell model experiments8,9, then the plausibility of past adaptation can be evaluated with the comparative method - that is, by asking whether a phenotype evolves repeatedly and independently under similar environmental pressures¹⁰ — or by using other, similarly indirect, approaches.

By contrast, analyses of 'time-stamped' ancient DNA data can provide more direct evidence of past human adaptation by facilitating precise 'evolution in action' tracking of genetic variants before, during and following

selection events^{11,12}. Excitingly, recent major advances in ancient DNA and sequencing methods (BOX 1) have facilitated a shift in this field from the analysis of mitochondrial genomes for a small number of individuals to the generation of genome-wide data sets (whole nuclear genome, exome, or genome-wide single-nucleotide polymorphism (SNP) data) at the scale of large populations^{13,14}. In fact, ancient genomes are now available for >1,100 archaic hominins and anatomically modern humans (FIG. 1a) representing time periods from the Middle Pleistocene (430,000 years before present (BP)) to the early 20th century (FIG. 1b; TABLE 1). Importantly for our understanding of human evolution, analyses of these data sets can provide high-resolution snapshots of the adaptive histories of genetically mediated behaviour, metabolism, soft tissue and other non-skeletal phenotypes not preserved in the fossil or archaeological records.

In this Review, we discuss how the informational content of ancient DNA can be maximized to explore the history of human adaptation at unprecedented depth and precision. We highlight the scope of approaches that enable the detection of selection signatures from ancient genomes, and we use an anthropological lens to explore examples from three major research areas: first, insights from direct analyses of ancient human genomes into adaptations to environmental change and related to the transition from hunter-gatherer to farming lifestyles; second, inferences of adaptive introgression (and the inferred functional effects) of genetic material from archaic hominins to modern humans; and third, ancient DNA-based characterizations of

Box 1 | Fuelling the ancient genomic explosion

Before 2010, ancient genome data sets were composed primarily of small numbers of archaic hominin individuals (FIG. 1a) and non-primate mammals^{150–153}. This pattern can be explained in part by the intense early excitement about archaic hominin genome sequencing¹⁵⁴. Yet, perhaps also contributing to this field's history is the difficulty in distinguishing authentic ancient human DNA from contamination introduced, for example, from human excavators, museum staff or laboratory personnel. Although potential contamination was a challenge regardless — as illustrated by analyses of early Neandertal palaeogenomic data¹⁵⁵ — the process of identifying authentic ancient DNA is less onerous when not studying the maximally similar (to those people excavating and handling the material) ancient genomes of anatomically modern humans^{98,99}. More recent studies have increasingly applied methodological and bioinformatic strategies that have helped to obviate this issue and to increase the confidence in ancient genomic data sets generated from anatomically modern humans¹⁵⁶. For example, modern contaminants can be distinguished from authentic ancient molecules based on characteristic damage patterns of cytosine deamination at sequence fragment ends in ancient DNA^{27,51}. Some laboratories now also perform a deamination repair step with a subset of each sample and then sequence both the unrepaired and repaired fractions in order to confirm authenticity while minimizing data loss and error^{69,157}.

Another trend in ancient human genomic studies has been the increasing reliance on relatively low-coverage sequence data (for example, $\leq 1x$ coverage, or an average of one or fewer sequence reads mapped to each nucleotide position of interest, per individual; FIG. 1a). This approach may also be combined with methods to enrich the sample for a targeted (but still substantial) fraction of the genome^{20,28} (TABLE 1). Depending on the analyses being performed, working with these data sets requires imputation of missing data by comparison with a modern human reference genome panel¹⁹, analytical focus on population allele frequency estimates rather than individual-level genotype calls²⁰, or other precautions⁶⁹. Such approaches greatly increase the feasibility of population-scale ancient DNA studies as a function of reduced sequencing costs (we note that as sequencing costs continue to decrease, we might eventually expect shifts towards higher proportions of more broadly powerful higher-coverage, whole-genome sequencing of ancient genome data sets). Combined with other laboratory methodological improvements^{27,158–160} and continuing sequencing technology advances¹⁶¹, these analytical innovations have helped fuel the recent rapid growth in the availability of ancient human genome data sets that may now be included in increasingly powerful evolutionary analyses.

> human-mediated selection on domesticated plant and animal species that provide insight into past human behaviour and evolution.

Positive natural selection

A mechanism of evolution in which a genetically mediated trait that confers a relative fitness advantage increases in frequency over time because of that advantage. In this Review, we refer to positive selection as an adaptive process that can act on new or previously existing genetic variants.

Phenotype

Physical traits of an organism; often refers to externally visible traits but may include internal and microscopic or biochemical traits.

Ancient DNA

DNA from palaeontological, archaeological, or historical but pre-modern biological specimens that is often damaged and degraded and recovered in small quantities. Tracking adaptation with ancient DNA

Positive natural selection is a nonrandom, directional process resulting in frequency increases of alleles associated with phenotypes that confer relative fitness advantages in a given spatiotemporal environmental context. Although this process has shaped patterns of modern human genetic diversity in ways that are detectable in present-day genomes^{6,15,16}, for most traits, it is only by turning to ancient DNA that we can establish precisely when and where such adaptive traits emerged.

The transformative power of ancient genomics for detecting and characterizing the genetic signatures of past human adaptations draws on our ability to access direct radiocarbon (¹⁴C) or indirect archaeological (for example, on the basis of pattern changes in tools, pottery or other artefacts associated with burials) estimates of the time period in which a particular individual lived. Armed with these data, researchers can explicitly track the trajectories of genetic variants before, during, and following selection events^{11,12,17,18}. Furthermore, ancient genome data can also be used to identify oscillating

allele frequency patterns that would otherwise be largely obscured from modern genome records^{19,20}. Precise knowledge of the spatiotemporal history of human adaptation can in turn facilitate integrative analyses of the specific environmental and cultural contexts associated with adaptive events and the corresponding implications for models of past human health and fitness.

The general evolutionary ancient genomics approach is straightforward: identify genetic variants with frequency changes significantly greater over a given time period than expected under genetic drift alone and/or relative to the genome-wide distributions of variant frequency changes (FIG. 2). This approach can be used to refine and test longstanding evolutionary hypotheses²⁰⁻²⁴ or to develop novel ones on the basis of newly uncovered signatures of selection²⁵. These strategies can also be extended to study polygenic adaptation, in which fairly subtle but consistent changes in the frequencies of multiple trait-associated loci can have large phenotypic effects in aggregate²⁶. Specifically, ancient genome data can be used to test whether shifts in the collective frequencies of a set of alleles known from a modern genome-wide association study (GWAS) to act directionally on a phenotype of interest — for example, stature — are unusual across a given time period compared with genome-wide backgrounds of frequency-matched alleles²⁰.

An important consideration to account for in evolutionary ancient genome analyses is that ancient human populations may not necessarily be closely related to subsequent or present-day people living in the same geographic regions. For example, recent ancient DNAbased studies have demonstrated a striking degree of repeated, large-scale human population movement and replacement across the past ~7,000 years in Europe^{19,27-30}, well beyond that previously predicted by even the most extreme archaeological models of population turnover³¹. Although such complex demographic histories can complicate ancient DNA-based interpretations of allele frequency change across time^{11,18}, methods to identify genome-wide and genome region-specific population structure can be incorporated into evolutionary ancient genome analyses to help distinguish between models of in situ change versus replacement or gene flow from a population with substantial differences in allele frequency^{20,32}.

Recent human evolution

The temporal distribution of available human ancient genomes (FIG. 1b) overlaps periods of major environmental and cultural change that are widely hypothesized to have represented significant drivers of recent human biological adaptation^{4,33}. For one, the expansion of anatomically modern humans to diverse habitats across the globe resulted in novel exposures to northern latitudes and cold climates³⁴. Two, the transition from hunting and gathering lifestyles to agriculture was associated with major dietary shifts³⁵, the construction of permanent settlements, and population density increases that in combination probably led to novel exposure to some pathogens and increases in the prevalence of others^{36,37}. Thus, evolutionary analyses of ancient genomes have



Exome

All or nearly all protein-coding gene regions of the nuclear genome; in humans, representing approximately 1% of the genome. Figure 1 | **The recent (and ongoing) ancient genomic explosion. a** | Cumulative counts of the numbers of ancient human and archaic hominin individuals with available ancient genomic data (whole genome sequences, exomes, and genome-scale single-nucleotide polymorphism data sets), by year of publication. For each category, both the total number of individuals and the subset of that sample with an average of >1 sequence read per targeted site are depicted. **b** | A spatiotemporal distribution of ancient human and archaic hominin genome data sets, where the areas of the circles are proportional to the genome counts. For some archaeological sites, precise geographic coordinates were not indicated in the corresponding publication; longitude and latitude were estimated in these cases. A summary of the published studies and data sets represented in the figure is provided in TABLE 1.

Reference	Year	Number of individuals with new ancient genome data*				Data type	Sample locations [‡]	Number of individuals per time period [§]			
		Archaic hominins		Ancient humans				>10,000 years BP	10,000- 5,000	<5,000- 2,500	<2,500 years BP
		Total	>1x	Total	>1x				years br	years br	
Noonan et al. ¹⁵⁰	2006	1	0			Direct cloning and WGS	Central Europe	1			
Green et al. ¹⁵¹	2006	1	0			PCR and 454 sequencing	Central Europe	1			
Green et al.98	2010	4	1			Restriction enzyme-based enrichment	Central, southern and eastern Europe	4			
Burbano et al. ¹⁶⁷	2010	0	1			SNP capture	Southern Europe	1			
Reich et al.99	2010	1	1			WGS	Northern and central Eurasia	1			
Rasmussen et al.156	2010			1	1	WGS	North America			1	
Rasmussen et al. ¹⁶⁸	2011			1	1	WGS	Oceania and Australia				1
Keller et al. ¹⁶⁹	2012			1	1	WGS	Southern Europe		1		
Skoglund et al.170	2012			4	0	WGS	Northern Europe		2	2	
Fu et al. ¹⁷¹	2013			1	1	Chromosome 21 capture	South and central Asia	1			
Castellano et al. ¹⁰⁸	2014	1	1			Exome	Central Europe	1			
Prufer et al. ¹⁰²	2014	1	1			WGS	Northern and central Eurasia	1			
Fu et al. ¹⁷²	2014			1	1	WGS	Northern and central Eurasia	1			
Schroeder et al. ¹⁷³	2014			3	0	WGS	North America				3
Gamba et al. ¹⁹	2014			13	8	WGS	Central Europe		9	4	
Lazaridis et al.52	2014			9	3	WGS	Central, northern and western Europe		9		
Malaspinas et al.174	2014			4	2	WGS	South America				4
Olalde et al.51	2014			1	1	WGS	Southern Europe		1		
Raghavan et al. ¹⁷⁵	2014			1	1	WGS	Northern and central Eurasia	1			
Rasmussen et al. ¹⁷⁶	2014			1	1	WGS	North America	1			
Seguin-Orlando et al. ¹⁷⁷	2014			1	1	WGS	Eastern Europe	1			
Skoglund et al. ¹⁷⁸	2014			7	2	WGS	Northern Europe		4	3	
Allentoft et al. ²⁷	2015			101	19	WGS	Central, northern, eastern and southern Europe; northern and central Eurasia		8	86	7
Fu et al. ¹⁷⁹	2015			1	1	Genome-wide SNPs	Eastern Europe	1			
Haak et al. ²⁸	2015			69	41	Genome-wide SNPs	Central, northern, eastern and southern Europe; northern and central Eurasia		45	24	
Mathieson et al.20	2015			83	31	Genome-wide SNPs	Central, eastern and southern Europe; Middle East; northern Eurasia		34	48	1

Table 1 | Human and archaic hominin ancient genome data sets

Reference	Year	Number of individuals with new ancient genome data*				Data type	Sample locations [‡]	Number of individuals per time period [§]			
		Archa homin	ic ins	Ancier humar	nt 1s			>10,000 years BP	10,000– 5,000 years BP	<5,000- 2,500 years BP	<2,500 years BP
Gallego Llorente	2015	Iotal	>1x	lotal	>1x 1	WGS	Africa			1	
Günther <i>et al.</i> ⁴⁷	2015			8	3	WGS	Southern Europe			8	
Jones et al. ⁴⁹	2015			3	3	WGS	Central and eastern Europe	2	1		
Olalde et al. ¹⁸¹	2015			6	1	WGS	Southern Europe		6		
Raghavan et al. ¹⁸²	2015			23	3	WGS	North and South America		2	1	20
Sawyer et al. ¹⁸³	2015	2	0			WGS	Northern Eurasia	2			
Rasmussen et al.184	2015			1	1	WGS	North America	1			
Cassidy et al. ³⁰	2016			4	3	WGS	Northern Europe		1	3	
Martiniano et al.185	2016			9	6	WGS	Northern Europe				9
Schiffels et al.186	2016			10	9	WGS	Northern Europe				10
Fu et al. ¹⁸⁷	2016			37	6	Genome-wide SNPs	Central, eastern, western and southern Europe; northern and central Eurasia	32	5		
Broushaki et al.50	2016			5	3	WGS	Middle East		4		1
Lazaridis et al.157	2016			44	16	Genome-wide SNPs	Middle East and eastern Europe	6	31	7	
Gallego Llorente et al. ¹⁸⁸	2016			1	1	WGS	Middle East		1		
Kılınç et al.189	2016			9	1	WGS	Middle East		9		
Jeong et al.54	2016			8	5	WGS	South and East Asia			1	7
Hofmanová et al. ⁶⁹	2016			7	5	WGS	Middle East and southern Europe		7		
Omrak et al. ¹⁹⁰	2016			2	0	WGS	Middle East		2		
Meyer et al. ¹⁹¹	2016	1	0			WGS	Southern Europe	1			
Lindo et al. ⁸¹	2016			24	25	Exome	North America			1	23
Skoglund et al. ¹⁹²	2016			4	4	WGS	Oceania and Australia			4	
Jones et al. ¹⁹³	2017			8	3	WGS	Northern and eastern Europe	1	7		
Unterländer et al. ¹⁹⁴	2017			8	0	Genome-wide SNPs and WGS	Northern and central Eurasia			4	4
Lipson et al. ¹⁹⁵	2017			127	26	Genome-wide SNPs	Central and southern Europe		93	34	
Lindo et al. ¹⁹⁶	2017			1	0	WGS	North America	1			
Kennett et al. ¹⁹⁷	2017			6	0	Genome-wide SNPs	North America				6
Mathieson et al. ¹⁹⁸	2017			195	95	Genome-wide SNPs	Central, eastern, Western, Northern, and Southern Europe	5	150	40	
Olalde et al. ²⁹	2017			165 [∥]	0	Genome-wide SNPs	Central, Northern, western and southern Europe		15	150	

Table 1 (cont.) | Human and archaic hominin ancient genome data sets

Reference	Year	Number of individuals with new ancient genome data*				Data type	Sample locations [‡]	Number of individuals per time period [§]			
		Archaic hominins		Ancient humans				>10,000 years BP	10,000- 5,000	<5,000- 2,500	<2,500 years BP
		Total	>1x	Total	>1x				years BP	years BP	
Martiniano et al.199	2017			14	7	WGS	Southern Europe		10	4	
Siska et al. ²⁰⁰	2017			5	0	WGS	Northern and Central Eurasia		5		
Haber et al.201	2017			5	4	WGS	Middle East			5	
González-Fortes et al. ⁴⁸	2017			6	6	WGS	Eastern and southern Europe		6		
Schuenemann et al. ²⁰²	2017			3	1	Genome-wide SNPs	Africa			2	1
Schlebusch et al. ²⁰³	2017			7	5	WGS	Africa				7
Mittnik et al. ²⁰⁴	2017			24	16	Genome-wide SNPs	Northern Europe, northern and central Eurasia		7	12	5
Slon et al. ²⁰⁵	2017	1	1			WGS	Northern and central Eurasia	1			
Günther et al.25	2017			7	4	WGS	Northern Europe		7		

Table 1 (cont.) | Human and archaic hominin ancient genome data sets

The authors apologize for any accidental omissions of published ancient human or archaic hominin genomic data sets. These data are otherwise intended to be current as of 25 July 2017. SNP, single-nucleotide polymorphism; WGS, whole-genome sequencing. *>1× indicates an average of greater than one sequence read mapped to each targeted site. If a new study generated additional DNA sequencing data for the same individual as included in a previous study, a new genomic data set was not recorded for that individual^{20,29,81,103,108,178,196,198,206}. However, for cases in which coverage was newly increased to >1×, this update was recorded^{20,81,167}. ⁴Geographic region definitions: central Europe: Croatia, the Czech Republic, Hungary, Austria, Poland, Germany, Switzerland, Serbia; eastern Europe: Belarus, the Ukraine, eastern Russia, Bulgaria, Montenegro, Caucasus, Romania; northern Europe: Estonia, Denmark, Sweden, Lithuania, Latvia, Ireland, United Kingdom, Norway, Netherlands, Belgium, France, Luxembourg; southern Europe: Spain, Portugal, Italy, Greece, Macedonia; South and East Asia: Nepal, China; northern and central Eurasia: Russia, Kazakhstar; Middle East: Iran, Israel, Turkey, Jordan, Lebanon; North America: Canada, United States, Greenland, Central America, the Caribbean; South America: Brazil, Chile; Africa: Ethiopia, Egypt. ⁴For small subsets of samples without reported specific dates^{27,45,50,81,181,190}, cultural periods were used to estimate time periods. IFor this study²⁹, sequence coverage per targeted SNP per sample was not provided. Among the samples, a range of 5,722–936,369 SNPs (of 1.24 million targeted SNPs) was covered by one or more sequencing reads. We recorded all samples as <1×; however, a small proportion of samples could actually cross this threshold.

Single-nucleotide polymorphism

(SNP). A single position in the reference genome at which the specific nucleotide present (thymine, guanine, cytosine, or adenine) varies among individuals in a population or species.

Archaic hominins

Now-extinct populations or species that are distinct from anatomically modern humans but that share a more recent common ancestor with modern humans than with chimpanzees — for example, Neandertals and Denisovans.

Anatomically modern humans

Hominins recognizable phenotypically as early members of our own species, *Homo sapiens*, first appearing > 200,000 years BP in Africa. the potential to confirm, refine, and expand our current understanding of the history of recent human biological adaptation.

Adaptation to novel environments. The roughly latitudinal gradient of ultraviolet A and B (UVA and UVB) radiation is thought to have been the principal evolutionary driver of modern human skin colour variation, with dark pigmentation in tropical latitudes protecting folate from photolysis, and depigmented skin in more northern and southern latitudes allowing for higher proportions of the fewer UVB rays to penetrate the skin to facilitate adequate vitamin D production³⁸. Vitamin D can also be acquired through a limited number of dietary sources, with fatty fish being one of the richest sources and smaller amounts found in beef liver, dairy products and egg yolk; however, this nutrient is largely absent from most agricultural plants³⁹⁻⁴¹. Vitamin D insufficiency is associated with cardiovascular disease, musculoskeletal disorders (for example, rickets), pre-eclampsia in pregnancy and other health concerns^{42,43}.

Given the predicted strong negative fitness effects of insufficient vitamin D production, it is interesting that recent ancient genome analyses have revealed a very gradual evolution of depigmentation among European

modern humans following their expansion into this region. Specifically, derived alleles associated with lighter skin pigmentation in the genes SLC45A2, SLC24A5, TYR and HERC2 are nearly fixed or are observed at their highest frequencies worldwide among living Europeans^{44,45}, with modern-human genome-based analyses estimating that selection acted on the majoreffect alleles SLC45A2 and SLC24A5 19,000-11,000 years BP⁴⁶. However, ancient genome data demonstrate that although the SLC45A2 allele was present as early as ~9,000 years BP in Scandinavia²⁵, it was absent or found at only fairly low frequencies at most sites across Europe before the onset of the Copper Age ~5,300-4,800 years BP^{19,20,27,47,48}. Meanwhile, although the *SLC24A5* allele was slightly more widely distributed by the Neolithic (that is, before ~7,000 years BP) in central and northern Europe^{19,20,25, 27,30,49} and in the Middle East⁵⁰, the ancestral allele was predominant in hunter-gatherers who lived ~13,000-7,000 years BP in what are now Spain (n = 3 individuals), Romania (n = 3), Switzerland (n = 1)and Luxembourg $(n = 1)^{48,49,51,52}$. Finally, *TYR* and HERC2 variants associated with lighter skin colour also experienced major frequency increases only fairly recently (for example, from 0.04 to 0.37 and from 0.16 to 0.65, respectively, between 6,500 and 4,000



genetic drift or purifying selection and are not observed to have changed substantially in frequency over the

frequency. **b** | Each variant across the genome is tested for an unexpectedly large temporal frequency shift.

investigated time period. Relatively rarely, a new mutation (or an existing mutation following environmental or

cannot be individually differentiated) arranged by human chromosome position on the x axis and negative log

cultural change resulting in different evolutionary pressures) confers a fitness benefit and may rapidly increase in

The statistical significance of each variant is shown as an individual point (due to the density of variants, most points

Adaptive introgression

The process of a genetic variant that was originally introduced into a population via admixture increasing in frequency by positive natural selection because it confers a fitness advantage.

P-value on the y axis.

mentioned above.

substantial European population turnover processes

observed a similarly staggered history of physiological

adaptations to the low-oxygen, high-altitude environ-

ment of the Himalayan Arc in Nepal (n = 5 individuals

from 3,150 to 1,250 years BP). Specifically, whereas

a derived ELGN1 allele associated with maintaining

oxygen homeostasis under high-altitude stress⁵⁵ was

represented in every sequence read mapped to this

position for all five individuals, derived EPAS1 vari-

ants associated with reduced haemoglobin levels that

In a South Asian ancient genome study, Jeong et al.54

Genetic drift

Changes in genetic variation over time that are due to random (chance) processes, apart from natural selection.

Gene flow

Movement of genetic variation between populations, for example, through migration or admixture.

Neolithic

A cultural period in human prehistory characterized by early technological and demographic shifts associated with the transition to farming and pastoralism, occurring at different times across regions. years BP in Ukraine)^{19,53}. Interestingly, in one study, the derived *HERC2* variant was observed at higher frequency in earlier hunter-gatherer samples than in later agriculturalist samples²⁰, perhaps reflecting the years BP; n = 2)⁵⁴.

> The gradual evolutionary histories of skin depigmentation and high-altitude-related adaptations among ancient Europeans and Himalayans, respectively, as revealed by ancient genomic analyses, may have several explanations. First, the random process of mutation is likely to play a part; that is, natural selection can only act on variation once it exists. Second, a time lag between the origin and geographic spread of an adaptive allele is to be expected^{57,58}, especially when considering the variability of ancient human migration patterns. Regardless, these ancient DNA-based evolutionary insights have implications for our understanding of prehistoric human

health and fitness. For example, does the finding of a relatively slow depigmentation process suggest that ancient Europeans were fairly susceptible to musculoskeletal or cardiovascular conditions associated with low levels of vitamin D? Alternatively, pre-agricultural Europeans could have ingested substantially more dietary vitamin D through hunting and fishing, as suggested by at least some archaeological evidence^{59,60}, to at least partially offset the expected reduction in skin-based production of this critical vitamin. Indeed, skeletal evidence of vitamin D insufficiency from the archaeological record is rare before the origins of agriculture, when it then begins to increase over time40. Future integrative studies that directly combine ancient genomic, archaeological and palaeopathological data sets might help researchers to more thoroughly characterize temporal variation in human health in the context of environmental change.

Biocultural adaptation to agricultural diets. Although some domestication processes probably had earlier roots^{61,62}, the archaeological record demonstrates that full transitions to agriculture for human societies were under way by ~12,000 years BP in the Middle East and not long after in Southeast Asia, the Americas and sub-Saharan Africa63. The subsequent near-worldwide spread of this technology led to novel evolutionary pressures associated with innumerable lifestyle, demographic and dietary changes. For example, one major dietary consequence of the agricultural transition was reduced nutritional diversity relative to the diets of many hunter-gatherer societies; starch alone comprises up to 70% of the caloric intake for some modern agriculturalist populations⁶⁴. Such specialization may have simultaneously resulted in new or increased evolutionary pressures on increasing digestive capacity for newly dominant food sources and maximizing metabolic efficiency for critical nutrients now deficient in some subsistence agricultural diets.

A classic example of biocultural adaptation is the co-evolution of the persistent production of the milk sugar lactose-digesting enzyme lactase into human adulthood with the cultural practice of collecting and consuming milk and dairy products from domesticated animals such as sheep, goats and cattle65. Without lactase persistence, human adults who consume lactose can have intestinal symptoms including diarrhoea, resulting in nutritional loss even beyond not receiving direct energetic benefits from lactose digestion⁶⁶. Human lactase persistence has evolved independently in multiple regions worldwide67. In Europe, the regulatory mutation conferring lactase persistence has been modelled based on modern human genetic data to have undergone selection beginning ~7,500 years BP⁶⁸, coincident with the spread of Neolithic (farming) cultures into this region. However, ancient DNA data clearly demonstrate a substantially more recent frequency increase and geographic spread of this adaptive variant only within the past ~5,000 years^{19,20,27,30,69,70} (FIG. 3; Supplementary table S1). Once again, this result has important implications for our models of prehistoric human health and behaviour, given that convincing archaeological evidence of dairy

production in Europe predates this new understanding of the emergence of the lactase persistence phenotype by more than two millennia^{71,72}. Before the spread of the adaptive allele, milk drinking behaviour by adults may have been more limited, and/or early dairy processes may have emphasized lactose-reducing processing techniques (for example, some cheese-making and fermentation processes), for which there is archaeological evidence from multiple European sites between 7,400 and 6,800 years BP^{73,74} (FIG. 3c).

Amylase, which is produced in humans by genes expressed in both the saliva (AMY1) and pancreas (AMY2A and AMY2B), is the enzyme responsible for the initial stages of starch digestion. Most present-day humans have an expanded number of AMY1 gene copies relative to chimpanzees, a difference associated with higher levels of salivary amylase protein expression⁷⁵. Human AMY1 duplications are associated with signatures of positive natural selection⁷⁶. The original AMY1 duplications post-date our divergence with the Neandertal and Denisovan lineage but pre-date modern human origins77. At least some modern human populations with traditionally high-starch diets have higher average AMY1 copy numbers than low-starch populations75, suggesting potential biocultural adaptation for expansion of the AMY1 locus following the agricultural transition. The few ancient DNA results reported to date are inconclusive with respect to this hypothesis; three European hunter-gatherers are estimated to have had five, six, and 13 diploid AMY1 gene copies⁵¹, whereas a Neolithic farmer from Germany (~7,000 years BP) had ~16 copies⁵². This test is an opportunity for future ancient genome analyses, even of existing data sets.

Current models of compensatory adaptations to nutritional deficiencies putatively associated with the transition to agriculture are also incomplete, but ancient DNA has an important role in our developing understanding of this evolutionary dynamic. For example, relative increases in the consumption of plants in some agricultural diets may have placed indirect evolutionary pressure on metabolic pathways of long-chain polyunsaturated fatty acids. These nutrients, which are consumed at higher levels in diets incorporating more animal protein (for example, from meat, fish or marine mammals), are critical for inflammatory responses and brain development⁷⁸. Human ancient genome data were used to identify a haplotype encompassing the fatty acid desaturase genes FADS1 and FADS2 that experienced marked frequency changes over the past ~4,000 years^{20,22,23}. Functional alleles in this haplotype can convert plant-based short-chain polyunsaturated fatty acids to long-chain fatty acids, thereby potentially compensating for the reduced dietary intake of these nutrients22.

Similarly, the amino acid ergothioneine, which is involved in protecting cells from oxidative stress and in the inflammatory response⁷⁹, must be obtained through dietary sources but is found at only very low quantities in domesticated cereal grains (for example, wheat and barley)⁸⁰. A functional variant in the ergothioneine transporter gene *SLC22A4* is hypothesized to have

Domestication

A process of plant and animal evolution mediated by human selection for particular phenotypes (artificial selection), sometimes combined with commensal adaptation to human-constructed niches.

Biocultural adaptation

The process of interaction between human cultural and adaptive biological change (for example, dairying and the ability of adults to digest milk sugars).

a Archaeological evidence of early Neolithic (7,500–5,000 BP) dairying in Europe



b Frequency of the European lactase persistence allele over time, by region



c Early archaeological evidence of lactose-reducing dairy processing techniques



Figure 3 | Biocultural adaptation to dairying and milk consumption. a | In addition to the early Neolithic presence of cattle and other domestic animals commonly milked for human consumption¹⁶³, early archaeological evidence of dairying practices in Europe include cattle-, sheep- and goat-specific β -lactoglobulin milk proteins recovered from human dental calculus and pottery fragments containing milk fat residues. b | Ancient DNA-based allele frequency estimates over time for the regulatory variant that confers the lactase persistence phenotype in Europe. For each population, the circle size represents ancient DNA sample size, and shading indicates geographic region. Centre time points reflect the midpoints of the date range estimates for each population sample. Allele frequencies estimated with PCR-based methods rather than from ancient genomic data sets are indicated with dashed circles; detecting and accounting for DNA damage and contamination is more difficult in PCR-based studies, potentially resulting in higher error rates. For reference, lactase persistence allele frequencies for present-day European populations are also indicated. A summary of the data used for this plot is provided in Supplementary table S1. c | Archaeological evidence of an early Neolithic dairy processing technique that would have resulted in reduced lactose contents for consumption by adults without lactase persistence. Shown is an ~7,200 years BP (before present) sieve fragment from Poland used for the production of lactose-reduced cheeses. Also shown is an ~3,800 years BP preserved piece of kefir (lactose-free) cheese from China made via fermentation. Part a (left panel) is reproduced with permission from REF. 162, CC-BY 4.0. Part a (middle panel) is reproduced with permission from REF. 164, CC-BY 4.0. Part a (right panel) is reproduced with permission from REF. 165, Macmillan Publishers Limited. Part c (left panel) image courtesy of Mélanie Roffet-Salque, University of Bristol, UK. Part c (right panel) is reproduced with permission from REF. 166, Elsevier.

provided protection against ergothioneine deficiency via increased absorption of this amino acid in European agriculturalists⁸⁰. However, although this allele was present at low frequencies in earlier stages of the Neolithic, ancient genome data have revealed that major frequency increases occurred only within the past 4,000 years²⁰. Thus, any farmers in this region who specialized in cultivating cereals earlier than 4,000 years BP would have been doing so without the widespread presence of this potentially compensatory allele. Alternatively, dietary diversity may have been greater in earlier time periods, limiting evolutionary pressure on the efficiency of this metabolic pathway until more recently.

Adaptation to a changing pathogen landscape. The

human burden of infectious diseases is hypothesized to have changed dramatically following the transition to agriculture^{36,37}. This phenomenon is thought to be explained by multiple factors, including the shift from small-group-based mobile hunting and gathering lifestyles to inhabiting permanent settlements with increasing population densities, being surrounded by human-modified habitats favoured by vector insects, and living in close proximity to livestock, which act as reservoirs of zoonotic pathogens. This period of novel exposure to some pathogens and increases in the prevalence of others was probably associated with intense new evolutionary pressure on the human immune system, including dynamic changes to the fitness spectrum (detrimental to neutral to beneficial) of functional alleles involved in pathogen defence.

Multiple ancient genome analyses have already helped to identify and characterize the trajectories of numerous adaptive alleles located within or near pathogen response and immune system genes in human populations worldwide^{20,50,51,69,81}. At present, equating particular ancient genome changes with evolved resistance to particular pathogens represents a major challenge due to both the broad immune system functions of many genes and uncertainty in the spatiotemporal distribution of exposure to multiple pathogens that were themselves also evolving. Yet, such inference may become increasingly possible with the improving power of modern disease-genotype association studies and the identification of subsets of alleles with relatively specific disease resistance effects, as for example with a number of Plasmodium falciparum and Plasmodium vivax malaria-resistance alleles²¹.

A different form of ancient genome data could ultimately provide major analytical benefits to these analyses by reducing the spatiotemporal uncertainty of past disease presence or absence and of disease prevalence. Specifically, beyond reconstructing endogenous genomes from ancient human skeletal remains, researchers can also characterize the ancient genomes of various pathogens that may have infected individuals during life⁸², as recently done for tuberculosis (*Mycobacterium tuberculosis*)⁸³, leprosy (*Mycobacterium leprae*)⁸⁴, plague (*Yersinia pestis*)⁸⁵, malaria (*P. falciparum*)^{86,87}, smallpox (Variola virus)⁸⁸, cholera (*Vibrio cholerae*)⁸⁹, *Staphylococcus saprophyticus*⁹⁰ and others^{82,91}.

It is also possible with ancient DNA data to reconstruct the virulence characteristics of pathogen strains as they evolved over time⁸⁹. For example, Rasmussen et al.92 sequenced the genomes of seven Eurasian Yersinia pestis strains from 5,000-2,800 years BP and analysed these data with those from other published ancient and modern Y. pestis strains. Interestingly, all strains from before 3,600 years BP were missing the *vmt* gene, which encodes a protein needed for the viability of the pathogen in the gut of its vector, the flea93. By contrast, nearly all Y. pestis strains from ~2,900 years BP to today contained an intact ymt gene copy, thus marking one of the key factors probably associated with the bubonic plague transmission cycle. Conversely, a ~660 years BP Y. pestis genome sequence recovered from London Black Death victims did not contain any derived nucleotide changes not also observed in present-day Y. pestis genomes, suggesting a potentially important role for concomitant human behaviour and ecology in the Black Death pandemic rather than genetic changes in the pathogen itself⁹⁴.

Continued expansion of these efforts will help to more precisely define the spatiotemporal distributions of ancient pathogens and to model their effects on host morbidity and mortality, knowledge that is in turn important for testing human immune system coevolution hypotheses with endogenous ancient genome data, potentially even on local scales⁹⁵.

Archaic adaptive introgression

A rich history of intrigue surrounds the potential genetic contributions of archaic hominins to present-day modern humans. Hypotheses on this topic have ranged from the *in situ* evolution of anatomically modern humans from archaic hominin ancestors such as Neandertals in multiple continental regions, requiring extensive long-term gene flow⁹⁶, to a single sub-Saharan African origin of modern humans followed by widespread worldwide dispersal and the complete replacement of remnant archaic hominin populations, with no interbreeding⁹⁷.

Excitingly, not only have archaic hominin 'palaeogenomic' data helped to provide a definitive answer to this question — the correct model is intermediate between the two hypotheses⁹⁸ — but this field has also uncovered evidence of an additional, previously unsuspected contemporary archaic hominin lineage, the Denisovans⁹⁹. We now know that modern humans did largely replace non-African archaic hominin populations after expanding across the globe, but not before multiple interbreeding and admixture events, which have left a legacy of an average of ~1–4% Neandertal DNA in the genomes of Eurasian and Native American humans living today^{98,100–102} and of up to ~6% Denisovan ancestry in the genomes of indigenous Australians, Melanesians and some other Southeast Asian populations^{99,103–107}.

As a brief aside (but one that is of major interest for palaeoanthropologists), the presence of Neandertal and Denisovan alleles in modern humans living today means that there are two genomics-based pathways to help reconstruct archaic hominin evolutionary biology: one, through direct analyses of archaic hominin

Zoonotic

The ability of a pathogen to be directly or indirectly transmitted to humans from animals sharing the same habitat.

Admixture

Interbreeding between previously isolated populations.

palaeogenome sequences for loci where function can be inferred or tested^{77,108-110}; two, and perhaps much more powerfully, through GWAS or functional analyses conducted in modern humans that characterize the phenotypic effects of Neandertal or Denisovan ancestry alleles^{111,112}. These analyses will become increasingly informative as future GWAS are performed on increasing numbers of non-disease phenotypes and if more studies are conducted in populations with appreciable proportions of Denisovan ancestry.



Admixture with Neandertals and Denisovans played an important part in the history of modern human biological adaptation. In particular, these archaic hominins and their ancestors inhabited Eurasia for hundreds of thousands of years before the migration of anatomically modern humans out of Africa, with assumed long-term histories of biological adaptation to their local environments. Modern humans did not have this long history of ecological adaptation to the diverse habitats of Eurasia, but the genetic legacy of archaic hominin admixture could have provided fitness benefit opportunities for carriers of Neandertal or Denisovan alleles associated with environment-specific adaptive phenotypes¹¹³. Indeed, numerous signatures of adaptive introgression can be found in the genomes of Eurasian populations living today, identified as regions of the genome with unexpectedly large proportions (>0.50 in multiple cases) of Neandertal- or Denisovan-ancestry haplotypes relative to the much lower background levels of genome-wide archaic hominin ancestry^{100,114,115} (FIG. 4).

The benefits of Neandertal admixture for Eurasian modern humans may have included the immediate acquisition of alleles associated with lighter skin pigmentation, potentially benefiting vitamin D production in northern-latitude habitats with relatively low UVB levels. Among the genomic regions with the strongest signatures of adaptive introgression from Neandertals are genes involved in skin development and pigmentation, including a *BNC2* gene haplotype now observed at 70% frequency in present-day European populations and a *POU2F3* haplotype now observed at >60% frequency in East Asian populations¹⁰⁰. Nonetheless, this process certainly did not reflect an evolutionary end point for depigmentation in Eurasians (as discussed above).

Figure 4 | Adaptive archaic introgression. An illustration of genetic admixture between archaic hominins (that is, Neandertals or Denisovans) and anatomically modern humans including an example of adaptive introgression that is ultimately detectable from patterns of variation in modern human genomes. a | Chromosomes (or chromosome segments) with ancestry originally from archaic hominins and anatomically modern humans are represented in blue and orange, respectively. For illustrative purposes, only one pair of autosomal chromosomes is depicted. Chromosomes experience recombination with each generation; thus, archaic hominin ancestry segments become reduced in size over time. The location of an archaic hominin variant conferring a fitness advantage is indicated with a star. Following admixture and introgression, this variant (and the surrounding chromosome segment not broken down by recombination) increases in frequency over time in humans via positive natural selection. **b** | Representation of the proportion of haplotypes with archaic hominin ancestry (y axis) at points along this chromosome (x axis) observed in the descendent human population living today. The proportion of archaic hominin ancestry in the region containing the adaptively introgressed variant is exceptional. In other regions of the genome depauperate of archaic hominin haplotypes, archaic hominin alleles may have been deleterious and removed by purifying selection¹⁰⁰.

Meanwhile, the *EPAS1* haplotype associated with high-altitude tolerance in Tibetan populations was found to reflect adaptive introgression from Denisovans¹⁰⁴, even though the genome-wide proportion of Denisovan ancestry is much higher in other modern human populations^{107,116}. Similarly, a haplotype containing the *WARS2* and *TBX15* genes that is fixed at 100% frequency in Greenland Inuit populations¹¹⁷ also originated in Denisovans¹⁰⁵. The protein encoded by *TBX15* is involved in the differentiation of brown and inducible (brite) adipocytes, which have critical roles in heat production during cold exposure¹¹⁸, suggesting a potential evolutionary mechanism for this adaptive introgression event in populations inhabiting cold climates.

Genes involved in pathogen defence and immune response are also significantly enriched within adaptively introgressed regions of modern human genomes^{119,120}, with specific cases including MHC¹²¹, OAS¹²² and TLR^{113,119} gene loci. Multiple non-mutually exclusive scenarios might explain the inferred fitness benefits of these haplotypes upon their introgression to modern humans. First, some of these alleles could have co-evolved with archaic hominin-specific pathogens ultimately transferred to modern humans during the period of direct contact between populations^{123,124}. Second, this result may reflect the general pathogen defence benefit of increased individual genetic diversity (heterozygosity) that is known for many immune loci¹²⁵. Third, some immune system variants that originated by random mutation in the Neandertal or Denisovan lineages may simply have been particularly effective relative to ancestral or human lineage-specific alleles at the same loci. Continued experimental characterization and computational prediction of the functional effects of introgressed alleles at immune system loci120,126,127 could help researchers both distinguish among these scenarios and advance our general understanding of how our history of adaptive introgression affects the health of human populations living today.

Genomic signatures of domestication

Beyond ancient humans and archaic hominins, ancient genome data sets are also rapidly emerging for many other species, including human-domesticated animals and plants¹²⁸⁻¹³¹. In our opinion, evolutionary ancient genome analyses of domesticated taxa offer multiple extremely valuable windows into the history of human behaviour and evolution (FIG. 5). Specifically, ancient genome analyses that characterize the evolutionary origins and spreads of various domestication-associated traits provide indirect insight into the human choices that favoured their selection across different cultural periods¹³². Such work also yields information on the nutritional return of investments in crops and livestock from farming and pastoralism at various points in history^{129,131}, providing valuable data for models of past human-environment interaction, health and demography. Finally, this line of research can help to identify how and when domestic species may have adapted commensally (that is, with traits not directly selected for by humans) to environmental and human niche habitats

they shared with ancient people^{133,134}, thereby simultaneously advancing our understanding of evolutionary pressures faced by past humans themselves.

Recent ancient genome results for domesticated species have highlighted substantial temporal spacing in the emergence of key traits. For example, the analysis of ancient maize cobs from Mexico and the southwestern USA revealed that maize from 5,310–4,400 years BP already carried multiple functional, derived variants found in present-day varieties; these variants are associated with the complex traits of seed storage, flowering time and exposed edible kernel (rather than a hard shell). However, other ancestral variants associated with ear shattering (whereby seeds do not remain attached when ripened) and some starch production phenotypes were simultaneously present^{131,135}.

Ancient genome analyses of domestic chickens in Europe have demonstrated that selection for variants associated with yellow skin colour, reduced fear of humans and increased egg production occurred only over the past 1,100–500 years^{129,136}, counter to previous inferences of ~6,000 years¹³⁷. Interestingly, consistent with the timing of the emergence of these adaptive traits as demonstrated by ancient genome analysis, archaeological and historical evidence supports the notion that chicken husbandry intensified in medieval Europe as a result of religious fasting practices during this period that limited the consumption of four-legged animals¹²⁹.

To date, more evolutionary ancient DNA analyses have been published on horses than on any other domestic species. This collective body of work is providing strong insight into artificial selection behaviour by humans. For example, Librado *et al.*¹³⁸ studied 13 ancient genomes of domestic horses from 2,700–2,300 years BP in the Kazakh Steppe in Asia, observing enrichment for genes expressed in mammary and pituitary tissues within genomic regions with signatures of positive selection. The authors suggest that this pattern might reflect a history of human-directed selection for phenotypes related to horse milking¹³⁸, a practice conducted for millennia in the region according to archaeological pottery lipid analysis and ethnographic records¹³⁹.

Based on knowledge of the relationship between genetics and coat colour and pattern variation in modern horses, it is possible to make inferences about prehistoric horse phenotypes by using ancient DNA. Ludwig et al.¹⁴⁰ reported a marked increase in the diversity of coat colours and patterns in European early domesticated horses (6,000–2,000 years BP; n = 73 individuals) relative to horses before the onset of domestication (>6,000 years BP; n = 16), with inferred black, white, chestnut, silver and spotted patterns in domesticated horses compared with bay and dun colourations in pre-domestic animals¹⁴⁰. Librado et al.¹³⁸ inferred similarly extensive coat colour diversity in the Kazakh Steppe horses, with cream, chestnut and spotted variants present in their sample, and a third study identified the increased frequency of a genetic variant associated with leopard-spotted coat patterns (from 0.06 to 0.35 in 16 pre-domestic and 83 domestic horses, respectively),



Figure 5 | **Insights into the timing of different trait evolution processes for domesticated species based on ancient DNA.** Domestic plant and animal ancient DNA studies can inform our understanding of past human behaviour and ecology. Shown are selected published examples for maize, horse, dog, cat, and chicken. The asterisk indicates that for some observable feature evolution examples, it has yet to be determined whether changes reflect human-directed selection or the relaxation of selective constraint related to changes in harvesting pressure from natural predators or other ecological processes (see text). As domestic plant and animal ancient genome data sets continue to expand, we expect increasing numbers of taxa to have multiple ancient-DNA-characterized traits that together intersect two or even all three of these categories of trait evolution processes.

despite this variant also being associated with nervousness and night blindness traits¹⁴¹. At least some of these changes were shown to reflect human-directed artificial selection as opposed to relaxation of functional constraint (which could have occurred, for example, via reduction in harvesting pressure from visual predators)¹⁴¹. In contrast to the coat changes associated with the early stages of domestication for horses, a recent ancient DNA study of African, European and Asian domestic cats (~8,500–100 years BP; n = 67 individuals) reported a fairly recent emergence of only ~700 years BP for the blotched (tabby) coat pattern now common in present-day domestic cats¹⁴².

Dogs have an even longer-term co-evolutionary relationship with humans than do horses or cats⁶¹. Results from evolutionary genomic analyses seem to reflect this history; for example, signatures of positive selection in dog genomes are enriched for genes involved in starch digestion¹⁴³, suggesting commensal biological adaptations to a human-like starch-rich diet. In fact, one selection signature surrounds the pancreatic amylase gene (AMY2B), for which dogs were found to have an average of ~15 diploid gene copies (n = 136 individuals) versus only two copies for all tested wolves $(n = 35)^{143}$. Subsequent studies have reported a greater average number of AMY2B copies in breeds with long-term associations with agriculturalists versus those associated with hunter-gatherer societies¹⁴⁴, suggesting remarkable potential convergence with human AMY1 copy number variation75. As with humans, dog ancient genomics studies would help to clarify the precise history of this adaptation. Early results have thus far been mixed; a maximum of three diploid AMY2B copies were observed among three total European Neolithic dogs (~7,000-5,000 years BP) in two studies^{145,146}, whereas another study reported a substantial average copy number increase beginning ~7,000 years BP (n = 13 Eurasian dogs from 15,000-4,000 years BP)133. Future studies and expanded spatiotemporal sampling will be necessary to fully characterize the timing and pace of dog AMY2B expansion and the potential relationship of this process to human cultural change.

Conclusions and future perspectives

We have illustrated the power of ancient genomics for studying processes of human biological adaptation. Potential insights from this approach complement, yet can also reach beyond, those from studies based on analyses of modern human genomes or based on the fossil and archaeological records alone. In particular, multiple high-precision temporal snapshots of human adaptation already provided by ancient genomics demonstrate substantial time lags between environmental or cultural changes and related biological adaptations - for example, skin depigmentation and lactase persistence in ancestral Europeans. Although these timelines are in some respects unsurprising given the random process of mutation, these results nonetheless have implications for our models of human health and fitness and hypotheses of compensatory cultural behaviours (FIG. 3). With increasing numbers of ancient genome data sets becoming available (FIG. 1), we anticipate expanded insights into the spatiotemporal dynamics of human adaptation, deeper analyses of the evolutionary histories of polygenic traits, and the transition from a primary focus on characterization of known human adaptations to the discovery of new ancient genomic signatures of past natural selection.

We note that existing ancient genome data sets are biased heavily towards Europe (especially) and parts of Asia (FIG. 1). Although this pattern partly reflects environment-based differences in ancient DNA preservation¹⁴⁷, disparities in global science funding and emphasis are also likely to play a part. Initial evolutionary ancient-genomic analyses on data sets from regions elsewhere have already delivered exciting insights^{54,81,148}. Concerted efforts to help close the geographical ancient genome data gap would lead to a more comprehensive picture of the history of human biological adaptation, including in response to a broader diversity of worldwide habitats and cultural and population history changes.

Finally, direct study of ancient human genomes is not the only useful approach for obtaining novel evolutionary anthropology insights using DNA. The domestication examples highlighted in this Review and the associated implications of these results for understanding past human behaviour are cases in point. As the generation of ancient genome data becomes increasingly feasible, we can look forward to dramatic increases in the taxonomic reach and spatiotemporal precision of domesticated plant and animal analyses and insights.

Moreover, the population and evolutionary histories of many non-domesticated, non-human species have been affected long term by human size-selective hunting and harvesting pressures, habitat modifications, accidental or purposeful translocations, and other behaviours¹⁴⁹. Evolutionary ancient genomic analyses of affected taxa could thus help to indirectly reconstruct the intensities of various human behaviours at different points in time as a valuable component of a broader, ecosystem-level view of human evolutionary history.

- Huxley, T. H. Evidence as to Man's Place in Nature 1. (Williams and Norgate, 1863). 2 Darwin, C. The Descent of Man. and Selection in
- Relation to Sex (John Murray, 1871). Fan, S., Hansen, M. E. B., Lo, Y. & Tishkoff, S. A. Going
- 3 global by adapting local: A review of recent human adaptation. Science 354, 54-59 (2016).
- 4 Laland, K. N., Odling-Smee, J. & Myles, S. How culture shaped the human genome: bringing genetics and the human sciences together. Nat. Rev. Genet. 11, 137-148 (2010).
- Scheinfeldt, L. B. & Tishkoff, S. A. Recent human adaptation: genomic approaches, interpretation and insights. Nat. Rev. Genet. 14, 692-702 (2013).
- 6. Sabeti, P. C. et al. Positive natural selection in the human lineage. Science 312, 1614–1620 (2006).
- 7. Raj, T. et al. Common risk alleles for inflammatory diseases are targets of recent positive selection. Am. J. Hum. Genet. 92, 517-529 (2013).
- Papandreou, I., Cairns, R. A., Fontana, L., Lim, A. L. & Denko, N. C. HIF-1 mediates adaptation to hypoxia by 8 actively downregulating mitochondrial oxygen consumption. Cell Metab. 3, 187-197 (2006)
- 9 Natarajan, V. T. et al. IFN-y signaling maintains skin pigmentation homeostasis through regulation of melanosome maturation. Proc. Natl Acad. Sci. USA 111, 2301-2306 (2014).
- 10. Norton, H. L. et al. Genetic evidence for the convergent evolution of light skin in Europeans and East Asians. Mol. Biol. Evol. 24, 710-722 (2006).
- 11. Pickrell, J. K. & Reich, D. Toward a new history and geography of human genes informed by ancient DNA. Trends Genet. **30**, 377–389 (2014).
- Nielsen, R. et al. Tracing the peopling of the world through genomics. Nature 541, 302-310 (2017).
- 13 Orlando, L., Gilbert, M. T. P. & Willerslev, E. Reconstructing ancient genomes and epigenomes. Nat. Rev. Genet. 16, 395-408 (2015).
- 14. Llamas, B., Willerslev, E. & Orlando, L. Human evolution: a tale from ancient genomes. Phil. Trans. R. Soc. B 372, 20150484 (2016).
- Nakagome, S. et al. Estimating the ages of selection signals from different epochs in human history. Mol. 15 Biol. Evol. 33, 657-669 (2016).
- Field, Y. et al. Detection of human adaptation during the past 2000 years. Science 354, 760-764 (2016) 17 Malaspinas, A.-S., Malaspinas, O., Evans, S. N. &
- Slatkin, M. Estimating allele age and selection coefficient from time-serial data. Genetics 192, 599-607 (2012).
- Sams, A. J., Hawks, J. & Keinan, A. The utility of ancient human DNA for improving allele age estimates, with implications for demographic models and tests of natural selection. J. Hum. Evol. 79. 64-72 (2015).
- 19. Gamba, C. et al. Genome flux and stasis in a five millennium transect of European prehistory. Nat. Commun. 5, 5257 (2014).
- 20 Mathieson, I. et al. Genome-wide patterns of selection in 230 ancient Eurasians. Nature 528, 499-503 (2015)
 - This paper performed a genome-wide scan for signatures of positive selection with ancient genome data from 230 Europeans (~8,500-2,300 vears BP) and characterized the spatiotemporal frequency trajectories of adaptive alleles related to diet, skin pigmentation, stature and the immune response.
- 21 Gelabert, P., Olalde, I., de-Dios, T., Civit, S. & Lalueza-Fox, C. Malaria was a weak selective force in ancient Europeans. Sci. Rep. 7, 1377 (2017).
- 22. Buckley, M. T. *et al.* Selection in Europeans on fatty acid desaturases associated with dietary changes. Mol. Biol. Evol. 34, 1307-1318 (2017).

- 23. Ye, K., Gao, F., Wang, D., Bar-Yosef, O. & Keinan, A. Dietary adaptation of FADS genes in Europe varied across time and geography. Nat. Ecol. Evol. 1, 0167 (2017).
- Sverrisdottir, O. Ó. et al. Direct estimates of natural 24 selection in Iberia indicate calcium absorption was not the only driver of lactase persistence in Europe. Mol. Biol. Evol. 31, 975-983 (2014).
- Günther, T. et al. Genomics of Mesolithic Scandinavia 25 reveal colonization routes and high-latitude adaptation. Preprint at bioRxiv http://dx.doi. org/10.1101/164400 (2017). This study conducted a genome-wide scan of positive selection using ancient genome data from seven Scandinavian individuals (~9,500-6,000 years BP) to
- reveal a haplotype that the authors propose may underlie physiological adaptation to cold climate. 26. Stephan, W. Signatures of positive selection: from
- selective sweeps at individual loci to subtle allele frequency changes in polygenic adaptation. Mol. Ecol. 25, 79-88 (2016).
- 27 Allentoft, M. E. et al. Population genomics of Bronze Age Eurasia. Nature 522, 167–172 (2015).
- 28 Haak, W. et al. Massive migration from the steppe was a source for Indo-European languages in Europe. Nature 522, 207-211 (2015).
- 29 Olalde, I. et al. The Beaker phenomenon and the genomic transformation of northwest Europe. Preprint at bioRxiv http://dx.doi.org/10.1101/135962 (2017).
- Cassidy, L. M. et al. Neolithic and Bronze Age 30. migration to Ireland and establishment of the insular Atlantic genome, Proc. Natl Acad. Sci. USA 113. 368-373 (2016).
- Shennan, S. Evolutionary demography and the population history of the European early Neolithic. Hum. Biol. 81, 339-355 (2009).
- Leonardi, M. et al. Evolutionary patterns and processes: 32 lessons from ancient DNA. Syst. Biol. 66, e1-e29 (2017).
- Mirazón Lahr, M. The shaping of human diversity: filters, boundaries and transitions. Phil. Trans. R. Soc. B 371, http://dx.doi.org/10.1098/rstb.2015.0241 (2016).
- 34 Stewart, J. R. & Stringer, C. B. Human evolution out of Africa: the role of refugia and climate change. Science 335, 1317-1321 (2012).
- Richards, M. A brief review of the archaeological evidence for Palaeolithic and Neolithic subsistence. Eur. J. Clin. Nutr. 56, 1262-1278 (2002).
- Perry, G. H. Parasites and human evolution. Evol. 36 Anthropol. 23, 218-228 (2014).
- Pearce-Duvet, J. M. C. The origin of human pathogens: evaluating the role of agriculture and domestic animals in the evolution of human disease. Biol. Rev. 81.369-382 (2006).
- 38 Jablonski, N. G. & Chaplin, G. Human skin pigmentation as an adaptation to UV radiation. Proc. Natl Acad. Sci. USA 107, 8962-8968 (2010).
- 39 Chaplin, G. & Jablonski, N. G. Vitamin D and the evolution of human depigmentation. Am. J. Phys. Anthropol. 139, 451-461 (2009).
- Brickley, M. B. *et al.* Ancient vitamin D deficiency: long-40. term trends. Curr. Anthropol. 58, 420-427 (2017).
- Ovesen, L., Brot, C. & Jakobsen, J. Food contents and 41. biological activity of 25-hydroxyvitamin D: a vitamin D metabolite to be reckoned with? Ann. Nutr. Metab. 47. 107-113 (2003).
- Bodnar, L. M. et al. Maternal vitamin D deficiency 42. increases the risk of preeclampsia. J. Clin. Endocrinol. Metab. 92, 3517-3522 (2007).
- 43 Wang, T. J. et al. Common genetic determinants of vitamin D insufficiency: a genome-wide association study. Lancet 376, 180-188 (2010).
- Lao, O., de Gruijter, J. M., van Duijn, K., Navarro, A. & 44. Kayser, M. Signatures of positive selection in genes associated with human skin pigmentation as revealed

from analyses of single nucleotide polymorphisms. Ann. Hum. Genet. 71, 354-369 (2007).

- 45 Sturm, R. A. et al. Human pigmentation genes under environmental selection. Genome Biol. 13, 248-263 (2012).
- 46. Beleza, S. et al. The timing of pigmentation lightening in Europeans. Mol. Biol. Evol. 30, 24-35 (2013).
- 47 Günther, T. et al. Ancient genomes link early farmers from Atapuerca in Spain to modern-day Basques Proc. Natl Acad. Sci. USA 112, 11917-11922 (2015).
- González-Fortes, G. et al. Paleogenomic evidence for 48 multi-generational mixing between Neolithic farmers and Mesolithic hunter-gatherers in the Lower Danube Basin. Curr. Biol. 27, 1801-1810.e10 (2017).
- 49. Jones, E. R. et al. Upper Palaeolithic genomes reveal deep roots of modern Eurasians. Nat. Commun. 6, 8912 (2015).
- Broushaki, F. et al. Early Neolithic genomes from the 50. eastern Fertile Crescent. Science 353, 499-503 (2016).
- 51 Olalde, I. et al. Derived immune and ancestral pigmentation alleles in a 7,000-year-old Mesolithic European. Nature 507, 225-228 (2014).
- Lazaridis, I. et al. Ancient human genomes suggest 52. three ancestral populations for present-day Europeans. Nature 513, 409-413 (2014).
- 53 Wilde S et al. Direct evidence for positive selection of skin, hair, and eye pigmentation in Europeans during the last 5,000 y. Proc. Natl Acad. Sci. USA 111, 4832-4837 (2014).

This study used ancient DNA data for alleles known to be involved in human pigmentation variation to identify a history of positive natural selection and estimate the strength of selection for each locus.

Jeong, C. et al. Long-term genetic stability and a highaltitude East Asian origin for the peoples of the high valleys of the Himalayan arc. Proc. Natl Acad. Sci. USA 113 7485-7490 (2016) This is an ancient genome study of eight individuals from Nepal (3,150-1,250 years BP) that found staggered appearances and frequency increases

for several genetic variants known from studies of modern regional populations to be associated with physiological adaptation to high altitude.

- Lorenzo, F. R. et al. A genetic mechanism for Tibetan high-55. altitude adaptation. Nat. Genet. 46, 951-956 (2014). Peng, Y. et al. Genetic variations in Tibetan 56.
- populations and high-altitude adaptation at the Himalayas Mol Biol Evol 28 1075-1081 (2011)
- Ralph, P. & Coop, G. Parallel adaptation: one or many 57. waves of advance of an advantageous allele? Genetics 186, 647-668 (2010).
- 58 Novembre, J., Galvani, A. P. & Slatkin, M. The geographic spread of the CCR5 A32 HIV-resistance allele PLoS Biol 3 e339 (2005)
- 59. Richards, M. P., Schulting, R. J. & Hedges, R. E. M. Archaeology: sharp shift in diet at onset of Neolithic. Nature 425, 366-366 (2003).
- 60. Chaplin, G. & Jablonski, N. G. The human environment and the vitamin D compromise: Scotland as a case study in human biocultural adaptation and disease susceptibility. Hum. Biol. 85, 529-552 (2013).
- Druzhkova, A. S. et al. Ancient DNA analysis affirms 61. the canid from Altai as a primitive dog. PLoS ONE 8, e57754 (2013).
- Snir, A. et al. The origin of cultivation and proto-62 weeds, long before Neolithic farming. PLoS ONE 10, e0131422 (2015).
- Boivin, N. L. et al. Ecological consequences of human 63 niche construction: examining long-term anthropogenic shaping of global species distributions. Proc. Natl Acad. Sci. USA 113, 6388-6396 (2016).
- Copeland, L., Blazek, J., Salman, H. & Tang, M. C. Form and functionality of starch. Food Hydrocoll. 23, 1527-1534 (2009).

- Gerbault, P. *et al.* Evolution of lactase persistence: an example of human niche construction. *Phil. Trans. R. Soc. B.* **366**, 863–877 (2011).
- Campbell, A. K., Waud, J. P. & Matthews, S. B. The molecular basis of lactose intolerance. *Sci. Prog.* 88, 157–202 (2005).
- Tishkoff, S. A. *et al.* Convergent adaptation of human lactase persistence in Africa and Europe. *Nat. Genet.* 39, 31–40 (2007).
- Itan, Y., Powell, A., Beaumont, M. A., Burger, J. & Thomas, M. G. The origins of lactase persistence in Europe. *PLoS Comput. Biol.* 5, e1000491 (2009).
- Hofmanová, Z. *et al.* Early farmers from across Europe directly descended from Neolithic Aegeans. *Proc. Natl Acad. Sci. USA* 113, 6886–6891 (2016).
- Burger, J., Kirchner, M., Bramanti, B., Haak, W. & Thomas, M. G. Absence of the lactase-persistenceassociated allele in early Neolithic Europeans. *Proc. Natl Acad. Sci. USA* 104, 3736–3741 (2007). This was the first ancient DNA-based study of the history of the European lactase persistence allele; this study reported that the allele was not present in eight early Neolithic individuals (~ 7,500 years в^P) from four geographic sites, suggesting that the ability of individuals to digest lactose across their lifetimes likely post-dated the origin and spread of European dairying practices.
 Craig, O. E. *et al.* Did the first farmers of central and
- Craig, O. E. *et al.* Did the first farmers of central and eastern Europe produce dairy foods? *Antiquity* **79**, 882–894 (2005).
- Copley, M. S. *et al.* Direct chemical evidence for widespread dairying in prehistoric Britain. *Proc. Natl Acad. Sci. USA* **100**, 1524–1529 (2003).
- Treuil, R. Dikili Tash, village préhistorique de Macédoine orientale. 1, Fouilles de Jean Deshayes (1961–1975), vol. 2, Bulletin de Correspondance Hellénique Supplément 37 (Ecole française d'Athènes, 2004).
- Salque, M. *et al.* Earliest evidence for cheese making in the sixth millennium BC in northern Europe. *Nature* 493, 522–525 (2013).
- Perry, G. H. *et al.* Diet and the evolution of human amylase gene copy number variation. *Nat. Genet.* **39**, 1256–1260 (2007).
- Inchley, C. E. *et al.* Selective sweep on human amylase genes postdates the split with Neanderthals. *Sci. Rep.* 6, 37198 (2016).
- Perry, G. H., Kistler, L., Kelaita, M. A. & Sams, A. J. Insights into hominin phenotypic and dietary evolution from ancient DNA sequence data. *J. Hum. Evol.* **79**, 55–63 (2015).
- Mathias, R. A. *et al.* Adaptive evolution of the *FADS* gene cluster within Africa. *PLoS ONE* 7, e44926 (2012).
- Paul, B. D. & Snyder, S. H. The unusual amino acid L-ergothioneine is a physiologic cytoprotectant. *Cell Death Differ.* 17, 1134–1140 (2010).
- Huff, C. D. *et al.* Crohn's disease and genetic hitchhiking at IBD5. *Mol. Biol. Evol.* **29**, 101–111 (2012).
- 81. Lindo, J. *et al.* A time transect of exomes from a Native American population before and after European contact. *Nat. Commun.* **7**, 13175 (2016). In this paper, the authors sequenced the exomes of 25 ancient First Nations individuals (~ 6,200-800 years æ) from British Columbia, Canada, to identify an *HLA-DQA1* gene haplotype with a substantial frequency difference compared with the Tsimshian descendant population living in the region today, potentially reflecting adaptation to disease outbreaks associated with European colonization (which post-dated the ancient DNA time series).
- Harkins, K. M. & Stone, A. C. Ancient pathogen genomics: insights into timing and adaptation. *J. Hum. Evol.* **79**, 137–149 (2015).
- Bos, K. I. *et al.* Pre-Columbian mycobacterial genomes reveal seals as a source of New World human tuberculosis. *Nature* 514, 494–497 (2014).
- Roffey, S. *et al.* Investigation of a medieval pilgrim burial excavated from the leprosarium of St Mary Magdalen Winchester. *PLoS Negl. Trop. Dis.* 11, e0005186 (2017).
- Spyrou, M. A. *et al.* Historical *Y. pestis* genomes reveal the European Black Death as the source of ancient and modern plague pandemics. *Cell Host Microbe* 19, 874–881 (2016).
- Marciniak, S. *et al. Plasmodium falciparum* malaria in 1 st – 2nd century CE southern Italy. *Curr. Biol.* 26, R1220–R1222 (2016).
- 87. Gelabert, P. *et al.* Mitochondrial DNA from the eradicated European *Plasmodium vivax* and *P. falciparum* from 70-year-old slides from the Ebro

Delta in Spain. *Proc. Natl Acad. Sci. USA* **113**, 11495–11500 (2016).

- Duggan, A. T. *et al.* 17 th century Variola virus reveals the recent history of smallpox. *Curr. Biol.* 26, 3407–3412 (2016).
- Devault, A. M. *et al.* Second-pandemic strain of *Vibrio cholerae* from the Philadelphia cholera outbreak of 1849. *N. Engl. J. Med.* **370**, 334–340 (2014).
- 90. Devault, A. M. *et al.* A molecular portrait of maternal sepsis from Byzantine Troy. *eLife* **6**, e20983 (2017).
- Warinner, C. *et al.* Pathogens and host immunity in the ancient human oral cavity. *Nat. Genet.* 46, 336–344 (2014).

This study reported ancient DNA results from dental calculus collected from four European individuals (~ 1,000–750 years se), including documentation of the presence of various pathogenic bacteria and producing direct evidence that pig, sheep, wheat and cruciferous vegetables were consumed as part of the diet.

 Rasmussen, S. *et al.* Early divergent strains of *Yersinia* pestis in Eurasia 5,000 years ago. *Cell* **163**, 571–582 (2015).
 This paper sequenced seven Eurasian plague

(Yersinia pestis) ancient genomes (\sim 5,000–2,800 years BP) and, among other findings, discovered that a gene encoding a protein necessary for Y. pestis viability in the flea gut was absent from genomes prior to \sim 3,600 years BP; the subsequent acquisition of this gene through horizontal transfer likely helped facilitate the bubonic plague transmission cycle. Hinnebusch, B. J. *et al.* Role of Yersinia murine toxin in

- Hinnebusch, B. J. et al. Role of Yersinia murine toxin in survival of Yersinia pestis in the midgut of the flea vector. Science 296, 733–735 (2002).
- Bos, K. I. *et al.* A draft genome of *Yersinia pestis* from victims of the Black Death. *Nature* **478**, 506–510 (2011).
- Hummel, S., Schmidt, D., Kremeyer, B., Herrmann, B. & Oppermann, M. Detection of the CCR5-Δ32 HIV resistance gene in Bronze Age skeletons. *Genes Immun.* 6, 371–374 (2005).
- Wolpoff, M. H., Thorne, A. G., Smith, F. H., Frayer, D. W. & Pope, G. G. in Origins of Anatomically Modern Humans (eds Nitecki, M. H. & Nitecki, D.) V.) 175–199 (Plenum Press, 1994).
- Tattersall, İ. Out of Africa: modern human origins special feature: human origins: out of Africa. *Proc. Natl Acad. Sci. USA* **106**, 16018–16021 (2009).
- Green, R. E. et al. A draft sequence of the Neandertal genome. Science 328, 710–722 (2010).
- Reich, D. *et al.* Genetic history of an archaic hominin group from Denisova Cave in Siberia. *Nature* 468, 1053–1060 (2010).
- Vernot, B. & Akey, J. M. Resurrecting surviving Neandertal lineages from modern human genomes. *Science* 343, 1017–1021 (2014).
- Dannemann, M., Pr
 üfer, K. & Kelso, J. Functional implications of Neandertal introgression in modern humans. *Genome Biol.* 18, 61–72 (2017).
- Prüfer, K. *et al.* The complete genome sequence of a Neanderthal from the Altai Mountains. *Nature* 505, 43–49 (2014).

 Meyer, M. *et al.* A high-coverage genome sequence from an archaic Denisovan individual. *Science* 338, 222–226 (2012).

104. Huerta-Sánchez, E. *et al.* Altitude adaptation in Tibet caused by introgression of Denisovan-like DNA. *Nature* 512, 194–197 (2014).
This paper demonstrated that a genetic haplotype surrounding the *EPAS1* gene that underlies a physiological adaptation to high altitude in modern Tibetans is the result of adaptive introgression from Denisovans or a related archaic hominin population.
105. Racimo, F. *et al.* Archaic adaptive introgression in

- TBX15/WARS2. *Mol. Biol. Evol.* **34**, 509–524 (2017). 106. Skoglund, P. & Jakobsson, M. Archaic human ancestry
- in East Asia. Proc. Natl Acad. Sci. USA 108, 18301–18306 (2011).
- Reich, D. *et al.* Denisova admixture and the first modern human dispersals into Southeast Asia and Oceania. *Am. J. Hum. Genet.* 89, 516–528 (2011).
- Castellano, S. *et al.* Patterns of coding variation in the complete exomes of three Neandertals. *Proc. Natl Acad. Sci. USA* 111, 6666–6671 (2014).
- Lalueza-Fox, C. *et al.* A melanocortin 1 receptor allele suggests varying pigmentation among Neanderthals. *Science* 318, 1453–1455 (2007).
- 110. Lalueza-Fox, C., Gigli, E., de la Rasilla, M., Fortea, J. & Rosas, A. Bitter taste perception in Neanderthals through the analysis of the *TAS2R38* gene. *Biol. Lett.* 5, 809–811 (2009).

- McCoy, R. C., Wakefield, J. & Akey, J. M. Impacts of Neanderthal-introgressed sequences on the landscape of human gene expression. *Cell* 168, 916–927 (2017).
- 112. Simonti, C. N. *et al.* The phenotypic legacy of admixture between modern humans and Neandertals. *Science* **351**, 737–741 (2016). This study used electronic health record phenotypes from a large sample of modern human patients to associate alleles originally introgressed from Neandertals with increased risk of depression, skin lesions associated with sun
- exposure (actinic keratosis), and other phenotypes.
 113. Gittelman, R. M. *et al.* Archaic hominin admixture facilitated adaptation to Out-of-Africa environments. *Curr. Biol.* 26, 3375–3382 (2016).
 This paper analysed 126 genomic regions containing strong signatures of adaptive introgression from archaic hominins identified in a sample of geographically diverse human populations; these loci are significantly enriched for genes involved in the immune response and also contain multiple genes with known roles in skin pigmentation.
- 114. Racimo, F., Sankararaman, S., Nielsen, R. & Huerta-Sánchez, E. Evidence for archaic adaptive introgression in humans. *Nat. Rev. Genet.* 16, 359–371 (2015).
- 115. Sankararaman, S. *et al.* The landscape of Neandertal ancestry in present-day humans. *Nature* **507**, 354–357 (2014).
- 116. Vernot, B. *et al.* Excavating Neandertal and Denisovan DNA from the genomes of Melanesian individuals. *Science* **352**, 235–239 (2016).
- 117. Furnagalli, M. *et al.* Greenlandic Inuit show genetic signatures of diet and climate adaptation. *Science* 349, 1343–1347 (2015).
- 118. Gburcik, V., Cawthorn, W. P., Nedergaard, J., Timmons, J. A. & Cannon, B. An essential role for Tbx15 in the differentiation of brown and 'brite' but not white adipocytes. *Am. J. Physiol. Endocrinol. Metab.* 303, E1053–E1060 (2012).
- Deschamps, M. *et al.* Genomic signatures of selective pressures and introgression from archaic hominins at human innate immunity genes. *Am. J. Hum. Genet.* **98**, 5–21 (2016).
- 120. Enard, D. & Petrov, D. A. RNA viruses drove adaptive introgressions between Neanderthals and modern humans. Preprint at *bioRxiv* <u>http://dx.doi.</u> org/10.1101/120477 (2017).
- 121. Abi-Rached, L. *et al.* The shaping of modern human immune systems by multiregional admixture with archaic humans. *Science* **334**, 89–94 (2011).
- 122. Sams, A. J. *et al.* Adaptively introgressed Neandertal haplotype at the OAS locus functionally impacts innate immune responses in humans. *Genome Biol.* **17**, 246–261 (2016).
- 123. Sullivan, A. P., de Manuel, M., Marques-Bonet, T. & Perry, G. H. An evolutionary medicine perspective on Neandertal extinction. *J. Hum. Evol.* **108**, 62–71 (2017).
- 124. Houldcroft, C. J. & Underdown, S. J. Neanderthal genomics suggests a pleistocene time frame for the first epidemiologic transition. *Am. J. Phys. Anthropol.* 160, 379–388 (2016).
- 125. Key, F. M., Teixeira, J. C., de Filippo, C. & Andrés, A. M. Advantageous diversity maintained by balancing selection in humans. *Curr. Opin. Genet. Dev.* 29, 45–51 (2014).
- Nédélec, Y. *et al.* Genetic ancestry and natural selection drive population differences in immune responses to pathogens. *Cell* 167, 657–669 (2016).
- 127. Quach, H. *et al.* Genetic adaptation and Neandertal admixture shaped the immune system of human populations. *Cell* **167**, 643–656.e17 (2016).
- Park, S. D. E. *et al.* Genome sequencing of the extinct Eurasian wild aurochs, *Bos primigenius*, illuminates the phylogeography and evolution of cattle. *Genome Biol.* 16, 234–249 (2015).
- 129. Loog, L. *et al.* Inferring allele frequency trajectories from ancient DNA indicates that selection on a chicken gene coincided with changes in medieval husbandry practices. *Mol. Biol. Evol.* <u>http://dx.doi.org/10.1093/</u> <u>molbev/msx142</u> (2017).

This study of domestic chickens connected the ancient DNA-informed timing of a significant change in frequency for an allele associated with increased egg production to concomitant increases in the intensity of chicken husbandry as documented by historical and archaeological records.

- MacHugh, D. E., Larson, G. & Orlando, L. Taming the past: ancient DNA and the study of animal domestication. *Annu. Rev. Anim. Biosci.* 5, 329–351 (2017).
- Ramos-Madrigal, J. *et al.* Genome sequence of a 5,310-year-old maize cob provides insights into the early stages of maize domestication. *Curr. Biol.* 26, 3195–3201 (2016).

This study identified a mix of ancestral and derived functional genetic variants in maize from Mexico ~ 5,300 years BP, highlighting the gradual temporal process of trait evolution in this domestic species.

- 132. Schubert, M. *et al.* Prehistoric genomes reveal the genetic foundation and cost of horse domestication. *Proc. Natl Acad. Sci. USA* **111**, E5661–E5669 (2014).
- 133. Ollivier, M. *et al. Amy2B* copy number variation reveals starch diet adaptations in ancient European dogs. *R. Soc. Open Sci.* **3**, 160449 (2016).
- 134. Librado, P. *et al.* Tracking the origins of Yakutian horses and the genetic basis for their fast adaptation to subarctic environments. *Proc. Natl Acad. Sci. USA* **112**, E6889–E6897 (2015).
- 135. Jaenicke-Després, V. *et al.* Early allelic selection in maize as revealed by ancient DNA. *Science* **302**, 1206–1208 (2003).
- 136. Flink, L. G. *et al.* Establishing the validity of domestication genes using DNA from ancient chickens. *Proc. Natl Acad. Sci. USA* **111**, 6184–6189 (2013).
- West, B. & Zhou, B.-X. Did chickens go North? New evidence for domestication. J. Archaeol. Sci. 15, 515–533 (1988).
- Librado, P. *et al.* Ancient genomic changes associated with domestication of the horse. *Science* 356, 442–445 (2017).
- 139. Outram, A. K. *et al.* The earliest horse harnessing and milking. *Science* **323**, 1332–1335 (2009).
- 140. Ludwig, A. *et al*. Coat colour variation at the beginning of horse domestication. *Science* **324**, 485 (2009).
- 141. Ludwig, A. *et al.* Twenty-five thousand years of fluctuating selection on leopard complex spotting and congenital night blindness in horses. *Phil. Trans. R. Soc. B.* **370**, 20130386 (2014).
- 142. Ottoni, C. *et al.* The palaeogenetics of cat dispersal in the ancient world. *Nat. Ecol. Evol.* **1**, 0139 (2017).
- 143. Axelsson, E. *et al.* The genomic signature of dog domestication reveals adaptation to a starch-rich diet. *Nature* **495**, 360–364 (2013).
- 144. Arendt, M., Cairns, K. M., Ballard, J. W. O., Savolainen, P. & Axelsson, E. Diet adaptation in dog reflects spread of prehistoric agriculture. *Heredity* 117, 301–306 (2016).
- Botigué, L. R. *et al.* Ancient European dog genomes reveal continuity since the Early Neolithic. *Nat. Commun.* 8, 16082 (2017).
- 146. Frantz, L. A. F. *et al.* Genomic and archaeological evidence suggest a dual origin of domestic dogs. *Science* **352**, 1228–1231 (2016).
- 147. Kistler, L., Ware, R., Smith, O., Collins, M. & Allaby, R. G. A new model for ancient DNA decay based on paleogenomic meta-analysis. *Nucleic Acids Res.* 45, 6310–6320 (2017).
- 148. Fehren-Schmitz, L. & Georges, L. Ancient DNA reveals selection acting on genes associated with hypoxia response in pre-Columbian Peruvian Highlanders in the last 8500 years. *Sci. Rep.* 6, 23485 (2016).
- 149. Sullivan, A. P., Bird, D. W. & Perry, G. H. Human behaviour as a long-term ecological driver of nonhuman evolution. *Nat. Ecol. Evol.* 1, 0065 (2017).
- Noonan, J. P. et al. Sequencing and analysis of Neanderthal genomic DNA. Science 314, 1113–1118 (2006).
- 151. Green, R. E. *et al.* Analysis of one million base pairs of Neanderthal DNA. *Nature* **444**, 330–336 (2006).
- 152. Ramírez, O. *et al.* Paleogenomics in a temperate environment: shotgun sequencing from an extinct Mediterranean Caprine. *PLoS ONE* 4, e5670 (2009).
- Miller, W. *et al.* Sequencing the nuclear genome of the extinct woolly mammoth. *Nature* **456**, 387–390 (2008).
- Lambert, D. M. & Millar, C. D. Evolutionary biology: ancient genomics is born. *Nature* 444, 275–276 (2006).
- 155. Wall, J. D. & Kim, S. K. Inconsistencies in Neanderthal genomic DNA sequences. *PLoS Genet.* 3, 1862–1866 (2007).
- Rasmussen, M. *et al.* Ancient human genome sequence of an extinct Palaeo-Eskimo. *Nature* 463, 757–762 (2010).

- Lazaridis, I. *et al.* Genomic insights into the origin of farming in the ancient Near East. *Nature* 536, 419–424 (2016).
- Pinhasi, R. *et al.* Optimal ancient DNA yields from the inner ear part of the human petrous bone. *PLoS ONE* **10**, e0129102 (2015).
- 159. Gansauge, M.-T. & Meyer, M. Single-stranded DNA library preparation for the sequencing of ancient or damaged DNA. *Nat. Protoc.* 8, 737–748 (2013).
- 160. Carpenter, M. L. *et al.* Pulling out the 1%: wholegenome capture for the targeted enrichment of ancient DNA sequencing libraries. *Am. J. Hum. Genet.* **93**, 852–864 (2013).
- Hofreiter, M. *et al.* The future of ancient DNA: technical advances and conceptual shifts. *BioEssays* 37, 284–293 (2015).
- 162. Gron, K. J. *et al.* Cattle management for dairying in Scandinavia's earliest Neolithic. *PLoS ONE* 10, e0131267 (2015).
- 163. Zeder, M. A. & Hesse, B. The initial domestication of goats (*Capra hircus*) in the Zagros mountains 10,000 years ago. *Science* **287**, 2254–2257 (2000).
- 164. Cramp, L. J. E. *et al.* Neolithic dairy farming at the extreme of agriculture in northern Europe. *Proc. Biol. Sci.* 281, 20140819 (2014).
- 165. Warinner, C. *et al.* Direct evidence of milk consumption from ancient human dental calculus. *Sci. Rep.* 4, 7104 (2015).
- 166. Yang, Y. *et al.* Proteomics evidence for kefir dairy in Early Bronze Age China. *J. Archaeol. Sci.* **45**, 178–186 (2014).
- Burbano, H. A. *et al.* Targeted investigation of the Neandertal genome by array-based sequence capture. *Science* 328, 723–725 (2010).
- 168. Rasmussen, M. *et al.* An Aboriginal Australian genome reveals separate human dispersals into Asia. *Science* **334**, 94–98 (2011).
- 169. Keller, A. et al. New insights into the Tyrolean Iceman's origin and phenotype as inferred by whole-genome sequencing. Nat. Commun. 3, 698 (2012).
- Skoglund, P. *et al.* Origins and genetic legacy of Neolithic farmers and hunter-gatherers in Europe. *Science* 336, 466–469 (2012).
- Fu, Q. *et al.* DNA analysis of an early modern human from Tianyuan Cave, China. *Proc. Natl Acad. Sci. USA* 110, 2223–2227 (2013).
- 172. Fu, Q. *et al.* Genome sequence of a 45,000-year-old modern human from western Siberia. *Nature* **514**, 445–449 (2014).
- 173. Schroeder, H. *et al.* Genome-wide ancestry of 17 th-century enslaved Africans from the Caribbean. *Proc. Natl Acad. Sci. USA* **112**, 3669–3673 (2015).
- Malaspinas, A.-S. *et al.* Two ancient human genomes reveal Polynesian ancestry among the indigenous Botocudos of Brazil. *Curr. Biol.* 24, R1035–R1037 (2014).
- 175. Raghavan, M. *et al.* Upper Palaeolithic Siberian genome reveals dual ancestry of Native Americans. *Nature* 505, 87–91 (2014).
- 176. Rasmussen, M. *et al.* The genome of a Late Pleistocene human from a Clovis burial site in western Montana. *Nature* **506**, 225–229 (2014).
- 177. Seguin-Orlando, A. *et al.* Genomic structure in Europeans dating back at least 36,200 years. *Science* 346, 1113–1118 (2014).
- Skoglund, P. *et al.* Genomic diversity and admixture differs for Stone-Age Scandinavian foragers and farmers. *Science* **344**, 747–750 (2014).
 Fu, Q. *et al.* An early modern human from Romania
- 179. Fu, Q. *et al*. An early modern human from Romania with a recent Neanderthal ancestor. *Nature* **524**, 216–219 (2015).
- Callego Llorente, M. *et al.* Ancient Ethiopian genome reveals extensive Eurasian admixture in Eastern Africa. *Science* **350**, 820–822 (2015).
- Olalde, I. *et al.* A common genetic origin for early farmers from Mediterranean Cardial and Central European LBK cultures. *Mol. Biol. Evol.* **32**, 3132–3142 (2015).
- 182. Raghavan, M. *et al.* Genomic evidence for the Pleistocene and recent population history of Native Americans. *Science* **349**, aab3884 (2015).
- Sawyer, S. *et al.* Nuclear and mitochondrial DNA sequences from two Denisovan individuals. *Proc. Natl Acad. Sci.* USA **112**, 15696–15700 (2015).
 Raymussen. M. *et al.* The ancestry and affiliations of
- Rasmussen, M. *et al.* The ancestry and affiliations of Kennewick Man. *Nature* **523**, 455–458 (2015).
 Martiniano, R. *et al.* Genomic signals of migration and continuity in Patients Informed April 2 August 2015.
- continuity in Britain before the Anglo-Saxons. *Nat. Commun.* **7**, 10326 (2016).

- 186. Schiffels, S. *et al.* Iron Age and Anglo-Saxon genomes from East England reveal British migration history. *Nat. Commun.* 7, 10408 (2016).
- 187. Fu, Q. *et al.* The genetic history of Ice Age Europe. *Nature* **534**, 200–205 (2016).
- Callego Llorente, M. *et al.* The genetics of an early Neolithic pastoralist from the Zagros, Iran. *Sci. Rep.* 6, 31326 (2016).
- Kılınç, G. M. *et al.* The demographic development of the first farmers in Anatolia. *Curr. Biol.* 26, 2659–2666 (2016).
- Omrak, A. *et al.* Genomic evidence establishes Anatolia as the source of the European Neolithic gene pool. *Curr. Biol.* 26, 270–275 (2016).
- Meyer, M. *et al.* Nuclear DNA sequences from the Middle Pleistocene Sima de los Huesos hominins. *Nature* 531, 504–507 (2016).
- Skoglund, P. et al. Genomic insights into the peopling of the Southwest Pacific. Nature 538, 510–513 (2016).
- 193. Jones, E. R. *et al.* The Neolithic transition in the Baltic was not driven by admixture with early European farmers. *Curr. Biol.* 27, 576–582 (2017).
- Unterländer, M. *et al.* Ancestry and demography and descendants of Iron Age nomads of the Eurasian Steppe. *Nat. Commun.* 8, 14615 (2017).
 Lipson, M. *et al.* Parallel ancient genomic transects
- 195. Lipson, M. et al. Parallel ancient genomic transects reveal complex population history of early European farmers. Preprint at bioRxiv <u>http://dx.doi.</u> org/10.1101/114488 (2017).
- 196. Lindo, J. et al. Ancient individuals from the North American Northwest Coast reveal 10,000 years of regional genetic continuity. Proc. Natl Acad. Sci. USA 114, 4093–4098 (2017).
- 197. Kennett, D. J. *et al.* Archaeogenomic evidence reveals prehistoric matrilineal dynasty. *Nat. Commun.* 8, 14115 (2017).
- Mathieson, I. *et al.* The genomic history of southeastern Europe. Preprint at *bioRxiv* <u>http://dx.</u> <u>doi.org/10.1101/135616</u> (2017).
- 199. Martiniano, R. *et al.* The population genomics of archaeological transition in west Iberia: Investigation of ancient substructure using imputation and haplotype-based methods. *PLoS Genet.* **13**, e1006852 (2017).
- 200. Siska, V. et al. Genome-wide data from two early Neolithic East Asian individuals dating to 7700 years ago. Sci. Adv. 3, e1601877 (2017).
- Haber, M. *et al.* Continuity and admixture in the last five millennia of Levantine history from ancient Canaanite and present-day Lebanese genome sequences. *Am. J. Hum. Genet.* <u>http://dx.doi. org/10.1016/j.ajhg.2017.06.013</u> (2017).
- 202. Schuenemann, V. J. *et al.* Ancient Egyptian mummy genomes suggest an increase of Sub-Saharan African ancestry in post-Roman periods. *Nat. Commun.* 8, 15694 (2017).
- Schlebusch, C. M. *et al.* Ancient genomes from southern Africa pushes modern human divergence beyond 260,000 years ago. Preprint at *bioRxiv* <u>http://</u> dx.doi.org/10.1101/145409 (2017).
- Mittnik, A. *et al.* The genetic history of northern Europe. Preprint at *bioRxiv* <u>http://dx.doi.org/10.1101/113241</u> (2017).
- 205. Slon, V. *et al.* A fourth Denisovan individual. *Sci. Adv.* **3**, e1700186 (2017).
- Kuhlwilm, M. *et al.* Ancient gene flow from early modern humans into Eastern Neanderthals. *Nature* 530, 429–433 (2016).

Acknowledgements

The authors thank C. Bergey and R. George for discussion about the manuscript. This work was supported by grants from the National Science Foundation (BCS-1554834 and BCS-1317163; to C.H.P.).

Competing interests

The authors declare no competing interests.

Author contributions

The authors contributed equally to all aspects of this manuscript.

Publisher's note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

- SUPPLEMENTARY INFORMATION
- See online article: <u>S1</u> (table) ALL LINKS ARE ACTIVE IN THE ONLINE PDF